

The Q and QTc Study (Quinine and QTc Interval Prolongation Study)

Bacha Amin Khan, Wasil Khan, Amin Ullah and Muhammad Rehman

ABSTRACT

Objective: The objective of this study is to determine the effect of Quinine on QTc interval.

Study Design: Observational / descriptive study.

Place and Duration of Study: This study was conducted at the Medicine Department of Saidu Teaching Hospital Saidusharif Swat from 1st October 2015 to 1st October 2016.

Materials and Methods: A total of 100 patients both male and female who needed Quinine for febrile illness were included Pretreatment ECG and post treatment ECG (on the last day of Quinine treatment) was recorded. Patients were divided in three groups on the basis of pretreatment ECG

Group I: (50%) pretreatment ECG QTc interval was 361 to 400msec (mean) 380 msec.

Group II: (30%) pretreatment QTC interval 401 to 450msec with mean 425.5msec.

Group III: (20%) pretreatment QTC interval 451 to 485msec with mean 468msec.

Patients were given Quinine according to the protocol and post treatment QTc interval was recorded.

Results: Values were analyzed on student T- Test with a P value of 0.006.

Conclusion: It is concluded that Quinine increases the QTc interval to a significant level.

Key Words: QTc- corrected QT interval, t-test Student t – test

Citation of article: Khan BA, Khan W, Ullah A, Rehman M. The Q and QTc Study (Quinine and QTc Interval Prolongation Study). Med Forum 2016;27(12):56-49.

INTRODUCTION

Malaria is an important cause of morbidity and mortality. Affecting more than 1 billion people and causing 1 to 3 million deaths each year¹. In our country malaria behaves like epidemic because of unstable transmission². Majority of the complications are due to plasmodium falciparum³. Major problem in the management of malaria is chloroquine resistance. Resistant malaria can be managed with alternative drugs.⁵

Malaria with complications need parenteral Quinine, which inhibits the polymerization of the toxic heme molecule⁶. Falciparum malaria can cause multi organ dysfunction due to cytokines production and impairment of microcirculation.⁷ Complications can occur if there is high parasite load (> 5%).⁸

Although Quinine is effective for severe malaria but has a number of side effects, like prolongation of the QTc interval, which is a risk for arrhythmias. The QTc is calculated according to the Bazett's correction i.e. $QTc = QT / \sqrt{R-R}$ msec⁹.

Department of Medicine, Saidu Group of Teaching Hospital, Saidu Sharif, Swat

Correspondence: Dr. Bacha Amin Khan, Associate Professor of Medicine, Saidu Group of Teaching Hospital, Saidu Sharif, Swat.

Contact No: 0301-8067612

Email: bacha66@hotmail.com

MATERIALS AND METHODS

This study was carried out in the Medicine Department of Saidu Teaching Hospital Saidu sharif Swat over a period of 1 year from 1st October 2015 to 1st October 2016. 100 Patients were included in the study. 70% were males and 30% were females. The age was ranging from 13 to 65 years.

Fever was the consistent clinical presentation (100%). Other symptoms were found in different combination like, headache (90%), nausea and vomiting (80%), abdominal pain (30%), Jaundice (20%), loose motions (20%) and confusion (10%). The symptoms, signs and other data were recorded on proforma. In 50% patients malarial parasites were isolated while in 50% patients malarial parasites could not be isolated. Patients were started on Quinine according to the protocol.

Pretreatment ECG was recorded for QTc interval. Patients were divided in three groups on the basis of pretreatment ECG Group I (50%) pretreatment ECG QTc interval was 361 to 400msec (mean) 380 msec.

Group II (30%) pretreatment QTC interval 401 to 450msec with mean 425.5msec. Group III (20%) pretreatment QTC interval 451 to 485msec with mean 468msec.

Post treatment (last day of Quinine treatment) QTc was recorded again and then the results were compiled and tested on student t-test.

RESULTS

The results are shown in table 1 and table 2.

Table No.1:Detail comparison of pre-treatment and post treatment of QTc interval in patients.

Group	No. of Patients	Pre-treatment QTc interval			Post treatment of QTc Interval		
		Min	Max	Mean	Min	Max	Mean
I	50	361	400	380.5	413	450	431.5
II	30	401	450	425.5	451	485	468
III	20	451	485	468	501	560	530.5
Total	100						

Table No.2: Mean comparison of pre-treatment and post treatment of QTc interval in patients.

Group	No. of patients	Pre-Treatment QTc mean	Post-Treatment QTc mean
I	50	380.5	431.5
II	30	425.5	468
III	20	468	530.5

T-Test (P. value)=	0.00609659
--------------------	------------

DISCUSSION

Malaria is common and sometimes presents with different manifestations¹⁰⁻¹⁴. Because of the development of resistance¹⁵⁻¹⁷ alternative drugs are needed for the management. Falciparum malaria can be fatal especially in non-endemic areas¹⁸. For uncomplicated cases artemether and lumefantrine combination can be used¹⁹ although it is expensive for developing countries²⁰. Doxycycline can be combined with antimalarial drugs²¹. Quinine in its parenteral form is used for severe malaria, although its therapy is responsible for resistance²². Up to 50% resistance has been reported for Quinine in case of falciparum malaria, which has an annual incidence of 33%^{24,25} the resistance has developed because of helicases as a result of adaptation to the stresses of existence²⁶ some studies have shown better results with artesunate regarding parasite clearance and safety.^{27,28}

Quinine therapy needs closed monitoring especially if loading doses are considered for the management of patients²⁹⁻³¹. The QT interval which was first described by Wolff in 1950.³² Can be either prolonged congenitally (330, or because of the underlying heart diseases³⁴ or because of the drugs especially Quinine and Quinidine. Prolonged QTC interval has been considered increased risk factor for cardiovascular mortality³⁵. The, quinidine syncope, was described in 1964³⁶. Both quinine and quinidine prolong the QTc interval.

However simple prolongation does not make the patient prone to torsade de pointes³⁷. QTc is considered

prolonged if it is > 440msec in males and more than 450msec in females³⁸. Although there is no rigid consensus on the prolongation limits but QTc of more than 500msec is a risk for arrhythmia³⁹. In our study the range of pretreatment QTc was 380-388 and post treatment QTc was 431-530msec with mean QTc, variable for different groups. Although the prolongation in QTc interval is significant statistically with a P-value of 0.006 but only 20%

patients had gone to QTc interval of > 500msec in the post treatment phase. None of the patients developed arrhythmias. May be because of careful selection of patients, close monitoring and addressing the supportive care.

In view of this study we suggest that early diagnosis by optimal malarial test and microscopy^{40,41} and proper management will decrease mortality and morbidity and avoid emergence of resistance strains especially, if WHO guidelines are adopted⁴² which recommends 2 yearly monitoring for drug resistant strains.

Malaria will remain a major health problem until effective vaccines are developed^{43,44} and effective preventive measures are taken⁴⁵.

CONCLUSION

Quinine prolong the QTc interval but only 20% go to the interval of > 500msec. Even in this group arrhythmias were not documented.

Conflict of Interest: The study has no conflict of interest to declare by any author.

REFERENCES

- White NJ, Breman JG. Malaria and babesiosis. Diseases caused by red blood cell parasites. In: Kasper DL, Fauci NS, Longo DL, et al, editors. Harrison's principles of internal medicine. 16th ed. McGraw Hill; 2005.p.1218 -1232.
- Centre for disease control and prevention. Health information for international travel, 1999-2000, Atlanta, department of health and human services, 2000.
- World health organization: severe falciparum malaria Tran R Soc Trop Med Hyg 2000;94 (Suppl 1): 51.
- White NJ. The assessment of antimalarial drug's efficacy trends parasitol 2002;18: 865.
- Van VUGTM, et al. Treatment of uncomplicated multidrug resistant falciparum malaria with artesunate-proguanil. Clin Infect Dis 2002;35.
- Abramowicz M. Drugs for parasitic infections, Med litt Drugs ther 2000;40:1.
- Moss PJ, Irving WL, Anderson J. Infectious disease, tropical medicine and sexually transmitted infections. In: Kumar P, Clark M, editors. Kumar and Clarke's. 8th ed. Clin Med Saunders; 2012.p. 143-146.

8. Rosental PJ. Protozal and helminthic infections. In: papadakis MA, Mcphee SJ, editors. current medical diagnosis and treatment. McGraw Hill; 2016.p.1482-1523.
9. Meek S, Moris F. Introduction-II- basic terminology. In: Morris F, Bray WJ, Camm J, editors. ABC of clinical electrocardiography. 2nd ed. Blackwell; 2008.p.5-8.
10. Kochar DK, Agarwal P, Kochar SK, Jain R, Rawat N, Pokharna RK, et al. Hepatocyte 7 days function and hepatic encephalopathy in plasmodium falciparum malaria. QJm 2003; 96:505-12.
11. Krishna A, Karnad DK. Severe falciparum malaria, an important cause of multi organ failure in Indian intensive care unit patients. Crit Care Med 2003; 31; 2278-84.
12. Mahmood K, Jairamani KL, Abassi B, Mahar S, Samo AH, Talib et al. Falciparum malaria various presentations. PAK J Med Sci 2006;22: 234-37.
13. Christopher JM, Rowland M, Sanderson F, Mutabingwa TK. Clinical review science, Medicine and the future, malaria. BMJ 2002;325: 1221-4.
14. Rashid S, Barsuom. Malarial acute renal failure Am Neph J 2000; 11: 174-82.
15. Meman IA, Kanth N, Murtaza G. chloroquin resistance malaria in children. JPMA 1998; 48(4): 98-100.
16. Meman IA, Dominate malaria parasite species in hospitalized children in Nawab Shah and chloroquin resistance. Specialist Pak J Med Sci 1997;13(3): 245-8.
17. Arain A Shaikh, Shaikh A. Study of malaria in children at Liaquat medical college hospital. Pak Pediat J 1997; 21(2); 95-97.
18. Molyneux Malcolm. Malaria in non-endemic areas. Med Int 1997; 37(II) : 28-31.
19. Rasheed A, Mk Saba YA, Muhammad etal. Efficacy and safety of Artemether- lumefantrine in uncomplicated falciparum malaria in Liberia. J Pak Med Assoc 2011;61(2) : 131-134.
20. Laxminarayan R. Act now or later? Economics of malaria resistance. Am J Trop Med Hyg 2004; 71(suppl2): 187-95.
21. Ejaz A, Haq Nawaz K, Hussain Z, et al. Treatment of uncomplicated plasmodium malaria with quinine-doxycyline combination therapy. J Pak Med Assoc 2007;57(10):502-505.
22. Bhalli MA, Samiullah. Falciparum malaria A review of 120 cases. JCPSP Pak 2001;11: 3003-3.
23. Jamal MM, Ara J, Ali N. Malaria in pediatrics age group; a study of 200 cases. Pak Armed forces Med J 2005; 55: 77-7.
24. Nizamani MA, Kalar NA, Khuhs IA. Burden of malaria in sindpakistan; A two year Surveillance report. J Liaquat Med Health Sci 2006; 5: 76-83.
25. Arain A, shaikh S, Shaikh A. study of malaria in children at liaquat medical college hospital. Pak Paed J 1992; 21: 95-7.
26. Tutejaetal R. Emerging functions of helicases in regularization of stress survival in malaria parasite plasmodium falciparum and their comparison with human host; parasitol international (2016), Hhp://dx.doi.org/10.1016/j-parint2016.08.010.
27. South East Asian quinine artesunate malaria trial (SEAQUMAT) artesunate versus quinine for treatment of severe falciparum malaria. A randomized trial www.thelancet.com Aug 27, 2005. Vol 366; 717-724.
28. Karunajeewa H, Lim C, Hung Ty, Illet KF, et al. Safety evaluation of fixed combination pipraquin plus dihydroartemisinin in Cambodian children and adults with malaria. Br J Clin Pharmacol 2003; 57(1) 93-99.
29. Tariq M, Saleem T, Hamidullah, Mehraj V, et al. Efficacy and safety of quinine loading dose in patients with severe falciparum malaria at a tertiary care hospital in Pakistan; J Pak Med Assoc 2011; 61(1); 27-31.
30. Alkadi HO. Antimalarial drug toxicity: A review chemotherapy 2007;53: 385-391.
31. White NJ. Cardio toxicity of antimalarial drugs. <http://infection.thlancet.com>. Vol7 Aug 2007.
32. Wolff-L. Electro cardiography. Fundamentals and clinical applications- Philadelphia. WB. Saunders company 1950.
33. Jarvell A, lange-nielsen F. congenital Deaf mutism, functional heart diseases with prolongation of QT interval and sudden death. Am heart J 1957; 54: 59-68.
34. Khambata RB, The QT interval in myocardial infarction. B Med J 1953;11:808-811.
35. Schouten EG, Dekker JM, Meppelink P et al. QT interval prolongation predicts cardiovascular mortality in an apparently healthy population. Circulation 1991;84(4):1516-1523.
36. Selzar A, Wray HW. Quinidine syncope. Paroxysmal ventricular fibrillation occurring during treatment of chronic atrial arrhythmias. Circulation 1964; 30: 17-26.
37. Darpo B. spectrum of drugs prolonging QT interval and the incidence of torsades pointes, Eur Heart J Suppl 2001; 3 (suppl K): K 70-80.
38. KT Chang, et al. Association between C reactive protein corrected QT interval and presence of QT prolongation in hypertensive patients: www.science-direct.com. Kaohsung journal of medical science 2014;30: 310-315.
39. Drew BJ, califf RM. Funk M, et al. Practice standards for electrocardiographic monitoring in hospital setting. An American heart association scientific statement from councils on cardiovascular Nursing, Clinical Cardiovascular

- diseases in young endorsed by international society of computerized electrocardiography and the American association of critical care nurses. *Circulation* 2004;110: 2721-46
40. Khan SA, Anwar M, Hussain S, Qureshi A H, Ahmad M, Afzal S. Comparison of optional malarial test with light microscopy for the diagnosis of malaria. Toronto critical care Medicine symposium (adult and pediatric). Oct 21-Oct 23- 2004.
 41. HaraniMs, Beg MA, Khaleeq L, Adil SA, Kakepoto GN, Khurshid M. Role of ICT malaria immunochromatographic test for rapid diagnosis of malaria. *J Pak Med Assoc* 2006;56(4):167-171.
 42. Global report on antimalarial efficacy and drug resistance 2000-2010 world health organization, Geneva, Switzerland.
 43. Racie TL, et al. progress with plasmodium falciparum sporozoite (Pfspz)- based malarial vaccine . *Vaccine* 2015;833:4752-7461.
 44. Stanisic, et al. development of cultured plasmodium falciparum blood – stage malarial cell banks for early phase in vivo clinical trial assessment of antimalarial drugs and vaccines. *Malaria J* 2015;143.
 45. Jamali S. Role of pyrethroids in control of malaria amongst refugee population. *J Pak Med Assoc* 2011; 61(5): 486-490.

Electronic Copy