Original Article

Wounds **Repairing Drugs**

Effects of Lactobacillus (Acidophilus) on Cutaneous Wound Repair with Respect to Neutrophil Count in Comparison with **Steroid**

1. Hina Khan 2. Muhammad Asif Sheikh 3. Mazhar ul Haque 4. Aneela Qureshi 5. Amin Fahim 6. Nazia Qamar 7. Adnan Anwar

1. Asstt. Prof. of Anatomy, Al-Tibri Medical College, Karachi 2. Principal & Assoc. Prof. Isra Institute of Rehabilitation Sciences Karachi 3. Asstt. Prof. of Anatomy, 4. Prof. of Pathology, 5. Asstt. Prof. of Pathology, 6. Lecturer of Pathology, 7. Lecturer of Physiology, Al-Tibri Medical College, Karachi

ABSTRACT

Objective: To evaluate the effects of lactobacillus (acidophilus) in repairing of wound with respect to neutrophil count in comparison with steroid therapy.

Study Design: Experimental study

Place and Duration of Study: This study was conducted in the Department of atomy of Al-Tibri Medical College Isra University Karachi Campus from June 2014 to October 2014.

Materials and Methods: Randomly selected Wistar albino rats were taken and their weight was between 100 - 150 gms. The rats were divided in three groups on the basis of topical application. Group I was treated with topical normal saline, Group II with topical lactobacillus(acidophilus), and Group II opical Steroid were applied once daily).Each group is further divided into four sub groups according to the day of sampling like group Ia (day 03), group Ib (day 07), group Ic (day 14) and group Id (day 21). Samples are taken on day 3,7,14 & 21 from each group and tissues were processed and stained to observe the neutrophil count that showed rapid healing processes. Data was analyzed on SPSS version 21 by applying ANOVA and post hoc tukey's test.

Results: Results compared the neutrophil count of Lactob illegroup with steroid group and control group. Mean \pm SD of numbers of Neutrophils / x400 in group IIa ras20500 \pm 0.577, in group IIb11.250 \pm 1.258, in group IIc5.7500 \pm 0.5000, group IId3.7500 \pm 0.9574and in group lawas54.25 \pm 2.872, group Ib41.250 \pm 2.629, group Ic23.500 \pm 3.000 and group Id18.750 \pm 0.957. The semicant value was in group IIa and group Ia(P<0.000), group IIb and group Ib(P<0.000), group IIc and group (P<0.000) and group IId and group Id (P<0.000). Mean \pm SD of numbers of Neutrophils / x400 in groups IIa as 20.500 ± 0.577 , in group IIb 11.250 ± 1.258 , in group IIc 5.7500 ±.0.5000, group IId 3.7500 \pm 0.9574 and group IIIa was54.500 \pm 1.290, group IIIb 42.000 \pm 1.414, group IIIc32.000 \pm 0.8165 and group IIId28.750 \pm 1.258. The significant value was in group IIa and group IIIa (P<1.000), group IIb and group IIIb (P<0.002), group IIc and group IIIc (P<0.000) and group IIId and group IIId (P<0.000). Conclusion: Lactobacillus street output anti-inflammatory effect against both therapeutic groups. Key Words: Wound, Lactobacklus, Estaneous, Neutrophil

Citation of article: Khun H, Sheikh MA, Haque M, Qureshi A, Fahim A, Qamar N, Anwar A. Effects of Lactobacillus (Acidophiles) on Cutaneous Wound Repair with Respect to Neutrophil Count in Comparison with Steroid. Med Forum 2015; 26(10):30-32.

INTRODUCTION

The primary role of the skin, it serves as a protective barrier from the environment¹. The damage to the cell, tissues, organs and viscera including the skin is usually restored by a complex process termed as wound healing. Normal pattern of wound healing comprises a sequence of coordinated overlapping procedures or phases that engage acute and chronic inflammation and cell division³. Lactobacillus species and bifidobac-

Correspondence: Dr. Hina Khan, Assistant Professor of Anatomy, Al-Tibri Medical College, Karachi Cell No.: 0331-3504341 E-mail: draminfahim@gmail.com

tarium species were reported as the most frequently used probiotic strains⁴. Lactobacillus species are referred to a group of lactic acid-producing organisms⁵. Lactobacillus acidophilus bacteria produce

exopolysaccharides (EPS) that can be associated to a cell's surface or can discharge into the environment⁶. The main factor contributing to the process of healing is extracellular medium which is mainly composed of various structural and functional proteins⁷. After the initial injury while there is formation of fibrin clot, there is extravasation of neutrophils at the site of injury. This is usually accomplished within 24hours at the margins of wound⁸. Many researchers have focused the role of Lactic acid in wound repair as it is being consumed in cosmetic products such as moisturizers, body lotions and exfoliants⁹. Anabolic hormones such as, glucocorticoides, cortisol, mineralocorticoides, prolactin are involved in the protein synthesis¹⁰. The process of wound repair is also enhanced by Dehydroepiandosteron having potential in wound healing¹¹.

MATERIALS AND METHODS

Design of Study: Experimental study

Setting: Anatomy department of Al- Tibri Medical College Isra University Karachi campus. Conducted after the approval of ethical committee of Isra University from June 2014-October 2014.

Sampling Technique: Randomized

Sample size: A total of 60 Wister Albino rats were divided into three groups on the basis of treatment given. Group I was treated with topical normal saline, Group II with topical lactobacillus (acidophilus), and Group III topical Steroid were applied once daily. Samples were taken on day 3,7,14 & 21 from each group and tissues were processed and stained to observe the neutrophil count that showed rapid healing processes. Each group is further divided into four sub groups as follows; group Ia (day 03), group Ib (day 07), group Ic (day 14) and group Id (day 21). Data was analyzed on SPSS version 21 by applying ANOVA and post hoc tukey's test.

Isolation of Lactobacilli: Species of lactobacillus were obtained from yogurt and strains were identified by the Microbiology department of PCSIR, Karachi and grown on (MRS) agar medium and incubated at 25° C for 48 hours. The solution contained 10^{10} to 10^{11} CFU ml bacteria that were collected every day after 48 hours. **Wound Formation**: "Wound" was crutted on the dorsal surface of rats which was open with full-thickness measuring about 1.5×1.5 cm². Tissue was processed and stained to observe the neurophil count. **Data analysis procedure:** Data was analyzed on SPSS version 21 by Applying ANOVA and post hoc tukey's test.

RESULTS

The results compare the neutrophil count among the three therapeutic groups. Basically compare the other with the group applied lactobacillus (B). Results compared the neutrophil count of Lactobacillus group with steroid group and control group. Mean \pm SD of numbers of neutrophils / x400 in group IIa was 20.500 \pm 0.577, in group IIb 11.250 \pm 1.258, in group IIc 5.7500 ± 0.5000 , group IId 3.7500 ± 0.9574 and in group Ia was 54.25 \pm 2.872, group Ib 41.250 \pm 2.629, group Ic 23.500 \pm 3.000 and group Id 18.750 \pm 0.957. The significant value was in group IIa and group Ia (P<0.000), group IIb and group Ib (P<0.000), group IIc and group Ic (P<0.000) and group IId and group Id (P<0.000). Mean ± SD of numbers of neutrophils / x400 in groups IIa was 20.500 ± 0.577 , in group IIb 11.250 ± 1.258 , in group IIc 5.7500 ± 0.5000 , group IId 3.7500 ± 0.9574 and group IIIa was 54.500 ± 1.290 , group IIIb 42.000 \pm 1.414, group IIIc 32.000 \pm 0.8165 and group IIId 28.750 \pm 1.218. The significant value was in group III 28.750 \pm 7.00. The significant value was in group IIa and group IIIa (P<1.000), group IIb and group IIIb (P<0.002), group IIc and group IIIc (P<0.000) and group II and group IIId (P<0.000) as shown in Figure 1a and figure number 1.1 and 1.3, that showed the decrease number of neutrophil in lactobac us group as compared with steroid given group.





Photomicrograph 1.1 shows decreased neutrophil count/400X (H&E stain) of Group II. H (hemorrhage), F(Fibroblast), C (collagen fibers), N (



Photomicrograph 1.2 shows increased neutrophil count/400X (H&E stain) of Group I. Ad (Adipose tissue), BV(blood vessels) and N (neutrophil)





Lactobacillus can easily be obtained by normal flora and from the food source of our daily life. It has got potential benefits if it is used for the accurate purposes. Few of the studies have also revealed the beneficial role of lactobacillus on the natural micro intestine¹². The induction of live strains of Lactobacillus acidophilus activated the cytokines which in turn are involved in the re-epithelization by production of cell components¹³. Marked reduce count of neutrophil can be seemed in lactobacillus treated group in accordance with some of researcher studies¹⁴. In accordance with Halper J; our study showed significant reduction in neutrophil count in lactobacillus treated group, that's shows the potency of lactobacillus species as a strong anti inflammatory agent. On day 3-4 increase numbers of neutrophil shows inflammatory phase, but this study reports on day 3 decreased in count of neutrophil showed antiinflammatory role of lactobacilli in comparison with others¹⁵.

CONCLUSION

On the bases of this study it evaluated that Lactobacilli are potent anti-inflammatory agent, as they showed decrease count of neutrophil in inflammatory phase of healing in comparison with other therapeutic groups.

Conflict of Interest: The study has no conflict of interest to declare by any author.

REFERENCES

- 1. Cordoso CA, Favoreto S, Oliveira LI, Vancin JC, Barban GB, Ferraz DB, et al. Olex acid modulation of the immune response in wound healing a new approach for skin repair. J. Immunol 2010; 216(3): 409-15.
- 2. Arbiser JL. Angiogenesis and the kin: a primer. J Am Acad Dermatol 1996; 34-186–97.
- Nilani P, Pranavi A, Duraitamy B, Damodaran P, Subhashini V, Elanto K. Formulation and evaluation of wound healing dermal patch. Afr. J. Pharm. Pharmacol 2011; 5(9): 1252-1257.

- Foligné B, Dewulf J, Breton J, Claisse O, Lonvaud-Funel A, Pot B. Properties of nonconventional lactic acid bacteria: Immunomodulation by Oenococcus oeni. Int J Food Microbiol 2010; 140(2): 136-145.
- Laws A, Guy, Marshall V. Biosynthesis, characterisation and design of bacterial exo polysaccharides from lactic acid bacteria. Biotechnol 2001;19(8): 597–625.
- 6. Burd A, Huang L. Carbohydrates and cutaneous wound healing. Carbo Chem 2008; 17: 253-274.
- Shivanada N, Marshal JR, Isitor G. Wound healing potential of ethanolic extract of Kalanchoe pinnate Lam. Leaf-A preliminary study. Ind J Exper Biol 2010; 48: 572-576.
- 8. Janis JE, Kwon RK, Lalonde DH. A Practical Guide to Wound Healing. Plast Reconstr Surg 2010; 125: 230-244.
- Smith WP. Epidermal and dermal effects of topical Lactic acid. J Am Acri Dermatol 1996;35,388-391.
- Lebeer S, Class JJ, Vanderleyden J. Antiinflammatory potential of probiotics: Lipoteichoi acid makes difference. Trends Microbiol 2012; 201-10.
- 11. Wray C. Mammen J, Hasselgren P. Catabolic response to stress and potential benefits of nutrition support. Nutrition 200;18:97.
- Wara MS, Kavitha BTVV, Srikanth JG, Velmani
 G. Probiotics as Potential Therapies in Human Gastrointestinal Health. Int J Adv Pharm Sci 2010; 1: 96-110.
- 13. Boirivant M, Strober W. The mechanism of action of probiotics. Gastroenterol 2007; 23: 679-692.
- Nasrabadi H, Ebrahimi T, Banadaki D, Kajousangi T, Zahedi. Study of cutaneous wound healing in rats treated with lactobacillus plantarum on days 1,3,7. Afri J Pharm Pharmacol 2011;5(21): 2395-2401.
- 15. Halper J, Leshin LS, Lewis SJ, Li WI. Wound healing and angiogenic properties of supernatants from Lactobacillus cultures. Exp Biol Med 2003; 228: 1329–1337.