

Correlation of BMI with Cholesterol and Sugar in Adults

1. Javed Akhtar Rathore 2. Zulifqar Ali Kango, 3. Mohammed Saleem

1. Asstt. Prof. of Medicine, AJK Medical College Muzaffaraabad Azad Kashmir 2 Medical Specialist, CMH Multan
3. Consultant Physician, DHQ Hospital, Kotli Azad Kashmir.

ABSTRACT

Objective: This study was aimed to find the correlation of BMI with cholesterol and sugar level in adult

Place and Duration of Study: This study was carried out in Department of Medicine, Combined Military Hospital (CMH) Quita from 2006 to 2009.

Study Design: Prospective observational cross sectional studies

Materials and Methods: Individuals with different ages, sex were selected as study population. The cholesterol and fasting blood sugar were measured according to standard protocol. Height in centimeter and weight in kilograms of each individual was recorded and BMI calculated as kg/m^2 . Physical examination was done for everybody. The SPSS-20 was used for statistical significant analysis. The frequencies of variable and correlation between BMI, heights, weight, sugar and cholesterol were comprehensively analyzed.

Results: A total of 2,174 adults, 1,947 (89.56%) male and 227 (10.44%) female were included in study. Age range was between 20 and 55years. The mean age was 38.47 ± 12.66 . Mean BMI was 23.57 ± 2.58 . Mean cholesterol was 4.57 ± 0.60 . Mean fasting blood sugar (FBS) was 4.67 ± 0.75 . Mean weight 70.32 ± 9.1 Mean height 172.73 ± 7.85 . The correlation analysis revealed that weight, fasting blood sugar (FBS) and cholesterol had positive correlation with BMI [correlation coefficient of 0.734 ($p < 0.000$), 0.167 ($p < 0.000$), 0.164 ($p < 0.000$) respectively and height had negative correlation with BMI [-0.123 ($p < 0.000$)].

Conclusion: BMI is positively correlated with weight, RBS and cholesterol. The effect of age, sex, exercise and current medical status, this correlation is reduced.

Key Words: Body Mass Index, blood sugar, Cholesterol, Obesity

INTRODUCTION

High BMI has been associated with great morbidity and mortality worldwide.¹⁻² The obesity is hazardous health problems. Overweight is defined as BMI of 25 to 29.9 kg/m^2 and obesity as a BMI of $>30 \text{kg/m}^2$. The mortality increases with increase prevalence of obesity.³ The overweight and Obesity have significant association with hypercholesterolemia and adults diabetes mellitus (DM)⁴⁻⁵. Lipid metabolism is also adversely affected in obesity. The prevalence of these risk factors substantially increases with increasing BMI. Overweight and obesity are also known to be independent risk factor for atherosclerotic cardiovascular risk disease in adults⁶. Increased body weight is a major risk factor for the metabolic syndrome as well as coronary heart disease (CAD). Many studies have demonstrated that individuals with metabolic syndrome are at high risk for subsequent development of non insulin dependent diabetes mellitus (NIDDM).⁷⁻⁹ The Higher BMI and impaired glucose tolerance is prevalent in children and adolescents. The relationship between BMI, glucose and lipids have been studied earlier¹¹⁻¹⁴. The aim of the present study was to investigate the relationship of BMI with atherosclerotic cardiovascular risk factors like serum cholesterol and glucose level in adult population.

MATERIALS AND METHODS

This study was aimed to find association of BMI with blood sugars and cholesterol in adults. The data was collected during 2006 and 2009 by the team of expert doctors. This cohort population studied was belonging to same professional groups and socioeconomic class. The data regarding the name, gender, marital status, nature of job, smoking, medical and drug history was obtained, entered and analyzed in SPSS 20. Height and body weight were measured by anthropometry. Body Mass Index (BMI) was determined as weight divided by height squared (Kg/m^2). BMI was divided into three groups 1: ≤ 24.9 , group 2: 25–29.9, and group 3: ≥ 30 as normal, overweight and obesity and assigned different categories. Fasting blood glucose and total cholesterol of each individual was measured. Frequency of variables calculated.

Quantitative data was expressed as Mean \pm SD and comparison between genders analyzed by independent sample *t*-test. The comparisons of BMI groups were accomplished by one way ANOVA. Pearson's correlation coefficient for continuous data between BMI, weight, cholesterol, sugar and heights two tailed taken as significant at $p < 0.01$ levels was used to show correlation between these variables.

RESULTS

A total of 2,174 individuals, 1,947 (89.56%) male and 227 (10.44%) female were examined. The mean age was 38.47 ± 12.66 . Mean BMI was 23.57 ± 2.58 . Mean cholesterol was 4.57 ± 0.60 . Mean fasting blood sugar (FBS) was 4.67 ± 0.75 . Mean weight 70.32 ± 9.1 Mean height 172.73 ± 7.85 . Comparison of these parameters between males and females showed that BMI, weight, sugar and cholesterol were significantly higher in male but height though higher but statistically insignificant (Table-1). Overweight individuals had significantly higher level of FBS and cholesterol than individuals of normal weight (Table -2). Our correlation analysis revealed that weight, fasting blood sugar (FBS) and cholesterol had positive correlation with BMI [correlation coefficient of 0.734 ($p < 0.000$), 0.167 ($p < 0.000$), 0.164 ($p < 0.000$) respectively and height had negative correlation with BMI [-0.123 ($p < 0.000$)]. Table 2.

Table No.1: Gender Distribution of Cardiovascular Factors (Height, Weight, FBS and Cholesterol)

Parameters	Male n=1947	Female n=227	p-value
BMI	23.66 ± 2.66	22.58 ± 3.38	0.000
Height	174.28 ± 6.88	159.50 ± 6.88	0.950
Weight	71.82 ± 7.84	57.49 ± 9.60	0.000
Sugar	4.75 ± 0.89	4.44 ± 0.75	0.035
Cholesterol	4.57 ± 0.61	4.51 ± 0.53	0.039

Independent sample t-test for comparing means

Table No.2: Correlation between BMI, Height, Weight and Cholesterol (Correlation Coefficient (p))

	BMI	Height	Weight	Sugar	Cholesterol
BMI	1				
Height	-0.123 (0.000)	1			
Weight	0.734 (0.000)	0.578 (0.000)	1		
Sugar	0.167 (0.000)	0.038 (0.076)	0.112 (0.000)	1	
Cholesterol	0.164 (0.000)	0.024 (0.272)	0.115 (0.000)	0.172 (0.000)	1

Correlation is significant at the 0.01 level (2-tailed)

DISCUSSION

Overweight and obesity lead many complications including diabetes mellitus, dyslipidemia and CAD¹⁵. Most of the studies on CAD risk factors have been done in developed country¹⁶⁻¹⁷. The higher BMI increases these CAD risk factors¹⁸. In our study high BMI groups have significant statistical association with overweight and Obesity. These results have been demonstrated with many previous studies¹⁹⁻²⁰. Our study showed cholesterol, and Fasting blood glucose level have

positive correlation with BMI as reported in developing and developed population^{21,22}. Differences have been reported in India and other studies^{23,24}. High BMI has predisposition to CVD in either sex²⁵. In our study correlation of mean cholesterol and fasting blood glucose with BMI is statistical significant.

Limitations in our study was not direct measurement of body fat by using CT and MRI²⁶. Others limitations were non utilities of parameters like physical activities²⁷, dietary habits²⁸, level of education with smoking²⁹, and not measuring of visceral fats by utilization of waist and waist hip ratio^{30,31}. Mean age in our study is actually representative of adult population. Our population comprising of healthy individual in contrary to general population with mean BMI in our study was below mean BMI reported earlier in our country³². In Asian high BMI has significant association with obesity³³⁻³⁴. High BMI has also correlation with overweight which is related to high food intake, genetic makeup and lack of exercise. As matter of fact our population has great risk of obesity as observed in world. As High BMI has major impact as atherosclerosis on health and associated with CVD and stroke. Therefore BMI should be routinely checked in clinical practice and epidemiological studies should be carried out to assess the atherosclerotic related disease burden in our society in order to make guidance for our population.

CONCLUSION

BMI is positively correlated with weight, RBS and cholesterol and is negatively correlation with weight.

REFERENCES

1. Prospective Studies Collaboration. Whitlock G, Lewington S, Sherliker P, Clarke R, Emberson J, Halsey J, et al. Body-mass index and cause-specific mortality in 900 000 adults: collaborative analyses of 57 prospective studies. *The Lancet* 2009; 373(9669): 1083–96
2. Ismail J, Jafar TH, Jafary FH, White F, Faruqui AM, Chaturvedi N. Risk factors for non-fatal myocardial infarction in young South Asian adults. *Heart* 2004;90:259-263.
3. Freedman DM, Ron E, Ballard-Barbash R, Doody MM, Linet MS. Body mass index and all-cause mortality in a nationwide US cohort. *Int J Obes (Lond)* 2006;30:822–9.
4. Ko GT, Chan JC, Cockram CS, Woo J: Prediction of hypertension, diabetes, dyslipidaemia or albuminuria using simple anthropometric indexes in Hong Kong Chinese. *Int J Obes Relat Metab Disord* 1999;23:1136-1142.
5. Chan JM, Rimm EB, Colditz GA, Stampfer MJ, Willett WC. Obesity, fat distribution, and weight gain as risk factor for clinical diabetes in men. *Diabetes care* 1994;17:961–9.

6. Hubert HB, Feinleib M, McNamara PM, Castelli WP. Obesity as an independent risk factor for cardiovascular disease; a 26 years follow-up for participants in the Framingham Heart Study. *Circulation* 1983;67:968–77.
7. Nguyen NT, Magno CP, Lane KT, Hinojora MW, Lane JS. Association of hypertension, diabetes, dyslipidemia, and metabolic syndrome with obesity: findings from the national health and nutrition examination survey, 1999 to 2004. *J Am Coll Surg* 2008;207:928–34.
8. Hanson RL, Imperatore G, Bennett PH, Knowler WC. Components of the “metabolic syndrome” and incidence of type 2 diabetes. *Diabetes* 2002;51:3120–7.
9. Deurenberg-Yap M, Chew SK, Lin VF, Tan BY, van Staveren WA, Deurenberg P: Relationships between indices of obesity and its co-morbidities in multi-ethnic Singapore. *Int J Obes Relat Metab Disord* 2001;25:1554–1562
10. Sinha R, Fisch G, Teague B, Tamborlane WV, Banyas B, Allen K, et al. Prevalence of impaired glucose tolerance <http://www.ayubmed.edu.pk/JAMC/PAST/22-4/Faheem.pdf> 77 among children and adolescents with marked obesity. *N Eng J Med* 2002;346:802–10.
11. Chan JM, Rimm EB, Colditz GA, Stampfer MJ, Willett WC. Obesity, fat distribution, and weight gain as risk factors for clinical diabetes in men. *Diabetes Care* 1994;17: 961–969.
12. Johnson RK, Appel LJ, Brands M, Howard BV, Lefevre M, et al. Dietary sugars intake, and cardiovascular health. A scientific statement from the American Heart Association *Circulation* 2009; 120: 1011–1020.
13. Johnson RJ, Segal MS, Sautin Y, Nakagawa T, Feig DI, et al. Potential role of sugar (fructose) in the epidemic of hypertension, obesity and the metabolic syndrome, diabetes, kidney disease, and cardiovascular disease. *Am J Clin Nutr* 2007;86: 899–906.
14. Lustig RH, Schmidt LA, Brindis CD. The toxic truth about sugar. *Nature* 2012;487: 27–29.
15. Villjoen A, Sinclair AJ. Diabetes and insulin resistance in older people. *Med Clin North America* 2011;95: 615–629. Voulgari C, Tentolouris N, Dilaveris P, Tousoulis D, Katsilambros N, et al. Increased heart failure risk in normal-weight people with metabolic syndrome compared with metabolically healthy obese individuals. *J Am Coll Cardiol* 2011;58: 1343–1350.
16. World Health Organization. Global database on body mass index; 2012.
17. Danaei G, Finucane MM, Lu Y, Singh GM, Cowan MJ, et al. National, regional, and global trends in fasting plasma glucose and diabetes prevalence since 1980: systematic analysis of health examination surveys and epidemiological studies with 370 country-years and 2.7 million participants. *Lancet* 2011;378: 31–40.
18. Yoon KH, Lee JH, Kim JW, Cho JH, Choi YH, et al. Epidemic obesity and type 2 diabetes in Asia. *Lancet* 2006;368:1681–1688.
19. Joshi P, Islam S, Pais P, Reddy S, Dorairaj P, Kazmi K, et al. Risk factors for early myocardial infarction in South Asians compared with individuals in other countries. *JAMA* 2007;297: 286–294.
20. Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection Evaluation And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III) *JAMA* 2001;285: 2486–2497.
21. Deurenberg-Yap M, Chew SK, Lin VF, Tan BY, van Staveren WA, Deurenberg P: Relationships between indices of obesity and its co-morbidities in multi-ethnic Singapore.
22. Bhopal R, Hayes L, White M, Unwin N, Harland J, Ayis S, Alberti G. Ethnic and socio-economic inequalities in coronary heart disease, diabetes and risk factors in Europeans and South Asians. *J Public Health Med* 2002;24:95–105.
23. Dadeja V, Misra A, Pandey RM, Devina G, Kumar G, Vikram NK. BMI does not accurately predict overweight in Asian Indians in northern India. *Br J Nutr* 2001;86:105–112.
24. Sendhu HS, Koley S, Sandhu KS. A study of correlation between lipid profile and BMI in patients with diabetes mellitus. *J Human Ecol* 2008;24:227–9.
25. Bakari AG, Onyemelukwe GC, Sani BG, Aliyu IS, Hassan SS, Aliyu TM. Relationship between random blood sugar and body mass index in an African population. *Int J Diabetes Metab* 2006; 14:144–5.
26. Van der Kooy K, Seideel JC. Techniques for the measurement of visceral fat: a practical guide *Int J Obese Relat Metab Disord* 1993;17:187–96.
27. Rastogi T, Vaz M, Spiegelman D, Reddy KS, Bharathi AV, Stampfer MJ, et al. Physical activity and risk of coronary heart disease in India. *Int J Epidemiol* 2004;33:759–767.
28. Panagiotakos DB, Pitsavos C, Chrysohoou C, Vlismas K, Skoumas Y, Paliou K, et al. Dietary habits mediate the relationship between socio-economic status and CVD factors among healthy adults: the ATTICA study. *Public Health Nutr* 2008;11:1342–1349.
29. Molarius A, Seidell JC. Differences in the association between smoking and relative body weight by level of education. *Int J Obese Relat Metab Disord* 1997;21:189–196.

30. Halkjaer J, Sorensen TI, Tjonneland A, Togo P, Holst C, Heitmann BL. Food and drinking patterns as predictors of 6-year BMI-adjusted changes in waist circumference. *Br Nutr* 2004;92:735-48.
31. Lemieux S, Prudhomme D, Bouchard C, Tremblay A, Despres JP. A single threshold value of waist girth identifies normal weight subjects with excess visceral adipose tissue. *Am J Clin Nutr* 1996;64:685-93.
32. Hussain S, Hussain I, Sana B, Waheed K, Qaisera S. Association of type 2 diabetes mellitus with biometric variable: A study in Sir Ganga Ram Hospital, Lahore. *Ann King Edward Med Uni* 2009;15(2):48-53.
33. Haerens L, Vereecken C, Maes L, De Bourdeaudhuij I. Relationship of physical activity and dietary habits with body mass index in the transition from childhood to adolescence: a 4-year longitudinal study. *Public Health Nutr* 2010;13:1722-8.
34. Nissinen K, Mikkila V, Mannisto S, Lahti-Koski M, Rasanen L, Viikari J, et al. Sweets and sugar-sweetened soft drink intake in childhood in relation to adult BMI and overweight. The Cardiovascular Risk in Young Finns Study. *Public Health Nutr* 2009;12:2018-26

Address for Corresponding Author:**Dr. Javed Akhtar Rathore,**

Assistant Professor of Medicine,

AJK Medical College, Muzaffarabad

Consultant Physician AKCMH/SKBZ Hospital

MuzaffarabaNear Neelum Walkway Lower Plate

Muzaffarabad A.K

E-mail: drjavedrathore111@yahoo.com,

Mobile: 0355-8106847, 03038109000

Electronic Copy