

# Correlation of C Reactive Protein Elevation with Hypertension in Tertiary Care Hospital

Muhammad Iqbal, Suhail Ahmed Almani and Shafak Nazia

## ABSTRACT

**Objective:** To determine the correlation between serum CRP levels and high blood pressure in hypertensive patients.

**Study Design:** Observational / descriptive study

**Place and Duration of Study:** This study was conducted at the Medicine OPD of LUMHS, Jamshoro from May 2016 to November 2016.

**Materials and Methods:** This study was conducted on 101. All the hypertensive patients were incorporated, after taking informed consent while, smokers, diabetics, patients having any infection, and any other co morbidity were excluded from the study to avoid bias in the results. Blood samples were taken for assessment of C reactive protein. Information was gathered on self-made proforma.

**Results:** Mean age of the participants was  $41 \pm 2.03$  years. Majority of the cases 60 (59.40%) were noted with age group of >40 years. 55 (54.77%) patients were male while 14 (13.86%) were female. Mean SBP was  $146.66 \pm 18.99$  mm Hg, mean diastolic blood pressure was  $96.50 \pm 10.59$  mm Hg and mean CRP was  $2.8 \pm 0.10$  mg/dL. 65.35% patients had elevated C reactive protein. Positive correlation was found between CRP and SBP R-value 0.36 and also positive correlation was noted between CRP and DBP r-value 0.38.

**Conclusion:** CRP elevation was positively correlated with hypertension, but not strongly correlated.

**Key Words** C reactive protein, hypertension, Co-relation

**Citation of article:** Iqbal M, Almani SA, Nazia S. Correlation of C Reactive Protein Elevation with Hypertension in Tertiary Care Hospital. Med Forum 2017;28(2):68-70.

## INTRODUCTION

Hypertension (HTN) is the typical, asymptomatic, promptly detectable and for the most part effortlessly treatable sickness that prompts to deadly difficulties if left untreated.<sup>1</sup> In the world over around 1 billion peoples having HTN “333 million in financially created and 639 million in the economically developing countries”.<sup>2</sup> It is most often is asymptomatic in nature which leads to dangerous complications in untreated cases. Around 7.6 million mortality and 92 million disabilities worldwide were attributable to hypertension in 2001. It is the common cause of atherosclerosis and various cardiovascular diseases CVDs like CHD, renal failure, CHF, ischemic stroke and the peripheral vascular disease.<sup>3</sup> HTN is the multi factorial attribute that outcomes from net impact of ecological and hereditary constituents. Elements that may add to HTN incorporate plenty salt in the diet or alcohol consumption, depression, aging, hereditary qualities physical idleness, rich fat saturated diets and the family history.

Inflammation considered as the commonest pathogenic element in the development of pulmonary arterial hypertension (PH) with several studies showing a significant association between systemic inflammatory markers and PH.<sup>4</sup> Inflammation is also a precursor for the atherosclerosis progression and systemic hypertension. CRP is the plasma protein, having “ $\leq 1 \text{mg/L}$ ” in normal cases with levels elevate 100 fold in the injuries, inflammation and infection’s response. CRP impulsive the “somatic C-polysaccharides of the pneumoniae by streptococcus bacteria and 1<sup>st</sup> acute phase of the protein to be labeled.<sup>5</sup> CRP primarily synthesize through hepatic response to “IL-6” and “IL-1  $\beta$ ”, and the risk valuation marker, and had good facts as the stable, with big half-life as nineteen hrs and demonstrated minor alterations in the levels in b/w fresh and the frozen forms which makes it as a good marker for diagnosis.<sup>2</sup> In 2001, Bautista et al, 1<sup>st</sup> time evaluate the CRP levels in in hypertensive cases and found it is the independent risk factor for hypertension development.<sup>6</sup> Elevated CRP can development the HTN by reduction of endothelium-dependent relaxation by decreasing nitric oxide in the endothelial cells, in outcome vasoconstriction and improved endothelin1 production.<sup>7</sup> CRP can also encourage the atherosclerosis by highly regulating angiotensin 1 receptor manifestation.<sup>8</sup> HTN, if not treated timely than in the cases may develop lethal complications and may result in morbidity and mortality. HTN is commonest

Department of Medicine, LUMHS Jamshoro.

Correspondence: Dr. Muhammad Iqbal, Associate Professor, Department of Medicine, LUMHS Jamshoro.

Contact No: 0333-2700192

Email: saedaarain@yahoo.com.com

Received: December 23, 2016; Accepted: January 20, 2017

health issue in population in developing nations and also major cause for CVD and it has been stated that HTN is the portion of inflammation and many researchers reported that increases of CRP in HTN cases. To best of our knowledge, we had conducted this study on relationship of CRP and hypertension in our setup.

**MATERIALS AND METHODS**

This observational descriptive study was containing on 101 cases attending medicine OPD of LUMHS over a time of 7 months; May 2016 to November 2016. All the hypertensive patients were incorporated after taking informed consent while, smokers, diabetics, patients having any infection, endocrine pathologies, Recent illness, previous history of IHS, MI, stroke, PVS and other vasculitis Chronic inflammatory diseases like Systemic Lupus Erythromatosis, and any other co morbidity were excluded from the study to avoid bias in the results. After taking thorough history and doing examination, blood was taken for estimation of C reactive protein. Information was gathered on self-made proforma. SPSS version 16 was used for analysis of results.

**RESULTS**

Total 101 hypertensive cases were selected. Mean age of the participants was 41+- 2.03. 11(10.49%)patients were with age group of < 30 years,30(29.70%)having age group of 30-40 years while 60(59.40%)were noted with age group of > 40 years. 55(54.77%) patients were male while 14(13.86%) were female. Table:1.

**Table No.1: Demographic characteristics of Patients n=101**

| Demographic characteristics | Number / percentages |
|-----------------------------|----------------------|
| <b>AGE</b>                  |                      |
| less than 30 years          | 11(10.49%)           |
| 30-40 years                 | 30(29.70%)           |
| More than 40 years          | 60(59.40%)           |
| <b>Gender</b>               |                      |
| Male                        | 55(54.45%)           |
| Female                      | 14(13.86%)           |

**Table No.2: Patients distribution according to Hypertension and CRP (n=101)**


| Blood pressure and CRP  | Mean±SD            |
|-------------------------|--------------------|
| Systolic blood pressure | 146.66±18.99 mm Hg |
| Diastolic BP            | 96.50±10.59 mm Hg  |
| CRP                     | 0.8±0.10mg/dL      |

Mean SBP was 146.66+18.99 mm Hg, mean DBP was 96.50+10.59 mm Hg and mean CRP was 0.8+0.10 mg/dL. Table: 2.


65.35% patients had elevated C reactive protein Fig: 1.

In this study positive correlation had found between CRP and SBP R-value 0.36 Fig:2


In our study positive correlation was found between CRP and DBP R-value 0.38 Fig:3



**Figure No.1. Patients distribution according to frequency of elevated CRP (n=101)**



**Figure No.2. Correlation between diastolic BP and CRP n=101**



**Figure No.3. Correlation between diastolic BP and CRP n=101**

## DISCUSSION

In present study assessed the correlation of high blood pressure and CRP and we found 65.35% hypertensive patients had increased c-reactive protein. Similar findings were seen in the study conducted by Dawris S.<sup>9</sup> On other hand Sesso and workers mentioned that the elevated level of the CRP is a risk of hypertension development. This correlation between elevated hs CRP and new-onset HTN led Sesso et al to suggested that HTN is the inflammatory disease.<sup>10</sup> The ATTICA study found higher levels of hsCRP, TNF- $\alpha$  and other inflammatory markers in pre-hypertensives. This communication was independent of the other co-existing causes for cardiovascular diseases indicating that pre-hypertension might be an inflammatory condition.<sup>11</sup> HTN may develop to several inflammatory stimulation at walls of vessel those encourage production of pro inflammatory cytokines like as the TNF- $\alpha$ , IL-6 and CRP as the protection against factors of the injuries. Several studies reported that inflammatory markers like as CRP is the particular determinant of endothelium containing vascular function in cases having CHD and this condition can also exist in the cases having HTN.<sup>2,13</sup>

CRP prevents the nitric oxide formation through endothelial cells that in turn encourage vasoconstriction, adhesion of the leukocytes, and activation of the platelets, oxidation and thrombosis. Furthermore elevated CRP may high regulate the receptors of the angiotensin and improve the expression of the activator inhibitor-1 of plasminogen through endothelial cells<sup>13</sup>. These both conditions may develop the BP and promote the atherogenesis. In our study 34.65% smoker cases were with elevated CRP level and CRP mean was  $0.8 \pm 0.10$  mg/dL. While in the study conducted by Dar MS et al<sup>2</sup> reported that mean hsCRP in the patients with hypertension is 3.26 mg/L on comparing with normal cases as 1.56 mg/L p-value 0.001. This mean level was very high as compare to our study, this may because in our study sample size is very short. Further Dar MS et al<sup>2</sup> reported that the significant difference of hs-CRP elevation was also seen in cases with shorter hypertensive duration as 1 year, as compare to those cases having hypertension history more than 5 years P-0.01.

Results of this study show that c reactive protein was positively correlated with both SBP and DBP.

## CONCLUSION

In the conclusion of this study the C reactive protein was positively correlated with hypertension, while no strong correlation was found, therefore further

randomized studies are require to conform the findings of this study.

**Conflict of Interest:** The study has no conflict of interest to declare by any author.

## REFERENCES

- Mishra CP, Kumar S. Risk Factors Of Hypertension In A Rural Area of Varanasi. *IJPSM* 2011;42;1;101-10.
- Dar MS, Pandith AA, Sameer AS, Sultan M, Yousuf A, Mudassar S. hs-CRP: A potential marker for hypertension in Kashmiri population. *IJCB* 2010;1;25(2):208-12.
- Harrison's Principles of Internal Medicine. 18th ed. Chapter 247:2042-59.
- Özpelit E, Akdeniz B, Özpelit ME, et al. Prognostic value of neutrophil-to-lymphocyte ratio in pulmonary arterial hypertension. *JIMR* 2015; 43;5;661-71.
- Hirschfield GM, Peppas MB. C-reactive protein and cardiovascular disease: new insights from an old molecule. *CJM* 2003; 1;96(11):793-807.
- Bautista LE, Lopez-Jaramillo, Patricio, et al. Is C Reactive Protein an independent risk factor for essential hypertension? *J Hypertension* 2001;19 (5):857-61.
- Verma S, Li SH, Badiwala MV, et al. Endothelin antagonism and interleukin-6 inhibition attenuate the proatherogenic effects of C-reactive protein. *Circulation* 2002;105;16;1890-96
- Wang CH, Li SH, Weisel RD, Fedak PW, Dumont AS, Szmítko P, et al. C-reactive protein upregulates angiotensin type 1 receptors in vascular smooth muscle. *Circulation* 2003; 107(13):1783-90.
- Dawri S, Padwal MK, Melinkeri R. Evaluation of high sensitivity C-reactive protein and serum lipid profile in prehypertension and essential hypertension. *NJIRM* 2014;5(1).
- Sesso HD, Buring JE, Rifai N, et al. C-Reactive Protein and risk of developing hypertension. *JAMA* 2003;290(22):2945-51
- Chrysohoou C, Pitsavos C, Demosthenes B, et al. The ATTICA study: Association between prehypertension status and inflammatory markers related to atherosclerotic disease. *AJH* 2004;17: 568-73.
- Andor M. The role of inflammation in endothelial dysfunction and progression of atherosclerosis in metabolic syndrome. *Timisoara Med J* 2005; 55(4):330-4.
- LI Jian-jun. Inflammation in hypertension: primary evidence. *Chin Med J* 2006;119:1215-21.