

# Determine the Outcome of Newborn in Post-Term Pregnancy

Outcome of Newborn in Post-Term Pregnancy

Ashba Anwer<sup>1</sup>, Nazia Tufail<sup>2</sup> and Asma Mudassir<sup>1</sup>

## ABSTRACT

**Objective:** To assess the outcome of newborn among pregnant females who deliver after prolonged pregnancy.

**Study Design:** Descriptive study

**Place and Duration of Study:** This study was conducted at the Department of Obstetrics & Gynaecology, Islam Central Hospital, Sialkot from October 2018 to March 2019.

**Materials and Methods:** One hundred patients with pregnancy of 42 weeks or above were included. Patients with medical disorders, intrauterine demise or pregnancy complications were excluded. Identification of parameters regarding poor neonatal outcome was done. Gestational age, parity, fetal movement, age, mode of delivery, ultrasound, admission cardiotocogram (CTG) and past prolonged pregnancy were recorded.

**Results:** There were 39 (39%) between 20-25 years of age. Multigravida was found in 60 (60%) and emergency caesarean section 65 (65%). Most of the babies 64% were admitted to neonatal intensive care unit (ICU). Complication of meconium aspiration syndrome was found 67 (67%). No fetal mortality was observed.

**Conclusion:** Pregnancy should be managed before 42 weeks of gestation and should not allow to went post-term due to high rate of neonatal mortality and morbidity.

**Key Words:** Neonate, Post-term pregnancy, Gestational age

**Citation of articles:** Anwer A, Tufail N, Mudassir A. Determine the Outcome of Newborn in Post-Term Pregnancy. Med Forum 2019;30(8):100-103.

## INTRODUCTION

Post-term pregnancy is defined as pregnancy more than 40 weeks of gestation (294-days) and above from last menstrual period (LMP) is defined as prolonged/post-term pregnancy.<sup>1</sup> There is approximately 3-12% reported frequency of post-term pregnancy.<sup>2</sup> However, the real incidence is probably less since most frequent reasons of prolonged pregnancy diagnosis is incorrect dating.<sup>3</sup> Risk factors includes genetic factors, male gender of fetus, prior post-term and primiparity for actual post-term pregnancy.<sup>4</sup> In the first two trimesters, obesity and consumption of fish is most recently describe factor.<sup>5</sup> To calculate the estimated due date (EDD) traditionally the last menstrual period (LMP) has been used, but due to use of this method, incorrectness exist in females who have irregular cycle, have first trimester bleeding or who have no recent hormone birth control methods.<sup>6</sup>

Therefore, when estimate gestational age regularity and length of cycle should be taken into account other than last menstrual period date. In early pregnancy, ultrasonographic dating can improve estimated due date (EDD) reliability.<sup>7</sup> However, it is essential to understand error margin reported at several times in each trimester. Prolonged/post-term pregnancy is related with increased incidence of prolonged labour and vacuum assisted birth or forceps (operative delivery). Due to a large size baby, patients are at risk of vaginal birth trauma.<sup>8</sup> In a post-term pregnancy cesarean delivery is likely twice due to the size of baby. Wound complications and infections and postpartum hemorrhage is also increased the risk factors for patients.<sup>9</sup> In a post-term pregnancy there are also risk for newborn and fetus. Fetus may stop gaining weight, volume of amniotic fluid decreases and the function of placenta decreases towards the end of pregnancy. If the baby is large than birth injury may also occur. There is also risk of meconium aspiration for those babies who born after 40 weeks.<sup>10</sup> Prolonged pregnancy management in absence of other complications is contentious. Females should be offered induction after forty one weeks as recommended by guidelines of Royal College of Obstetricians and Gynecologists.<sup>11</sup> From upto 40-weeks increased antenatal monitoring should be offered to those females who decline induction, containing twice cardiotocography and single deepest amniotic pool estimation through ultrasound. Less than 8cm pool depth indicated increase risk of intrapartum to fetus.<sup>12</sup> If use expectant management, labour should induce at beginning of 43<sup>rd</sup> week as recommended by some sources.<sup>13,14</sup>

<sup>1</sup>. Department of Obstetrics & Gynaecology, Islam Medical College / Hospital, Sialkot

<sup>2</sup>. Department of Obstetrics & Gynaecology, Khawaja Muhammad Safdar Medical College, Sialkot.

Correspondence: Dr. Ashba Anwer, Associate Professor of Obstetrics & Gynaecology, Islam Medical College / Hospital, Sialkot

Contact No: 03016103646

Email: drshahiddiyal@gmail.com

Received: June, 2019

Accepted: July, 2019

Printed: August, 2019

## MATERIALS AND METHODS

This descriptive case series study was conducted at Department of Obstetrics & Gynecology Department of Obstetrics & Gynaecology, Islam Teaching Hospital Sialkot from 1<sup>st</sup> October 2018 to 31<sup>st</sup> March 2019. A total of 100 un-booked patients above 40-weeks duration of pregnancy were included in this study, who were sure about the date of last menstrual period or had 1<sup>st</sup> trimester ultrasound report and with regular menstrual series. The patients with medical disorders, intrauterine demise or pregnancy complications were not were excluded from the study. After obtaining informed consent data was collected and gestational age, fetal movement, past prolonged pregnancy, age, parity, CTG admissions and ultrasound findings were studied. Factors were specifically noted who affecting poor perinatal outcome like birth weight, meconium aspiration, neonatal sepsis, respiratory distress syndrome, asphyxia, birth trauma, APGAR score, jaundice, admission and stay duration in neonatal intensive care unit. The collected data was analyzed through SPSS 20.

## RESULTS

Most of the patients 39 (39%) were 20-25 years old, 30 (30%) patients between the age group of 26-30 years, 18 (18%) were in 31-35 years of age group and 13 (13%) were above the age of 35 years. The numbers of primigravida 40 (40%) patients and multigravida patients were 60 (60%). All patients have longitudinal lie as confirmed by the ultrasonographic findings and transverse lie not presented by any patient. 3-3.5 kg estimated fetus weight in the majority 79 (79%) while only 21 (21%) have 3.6 kg to 4 kg weight. In 90 (90%) cephalic presentation was observed and breech presentation was found in only 10 (10%). 110-150 per minute fetal heart rate was found in 82 (82%) while less than 110 per minute fetal heart rate was found in 18 (18%). The delivered babies, males babies were 65 (65%) and females 35 (35%). In 5 (5%) develop birth trauma due to instrumental delivery. The babies who shifted to neonatal intensive care unit (NICU) were 64 (64%). Within three days most of the babies 51 (51%) were discharged and between 4 to 5 days 37 (37%) babies were discharged and 12 (12%) were discharged between 6 to 7 days (Table 1). According to the patients biophysical score, 10 (10%) had 6/10 score, 50 (50%) had 8/10 and 40 (40%) patients had 10/10 score (Table 2).

According to the delivery mode, most of the patients 65% had emergency C-section followed by spontaneous, instrumental and elective C-Section 20%, 12% and 3% respectively (Table 3). Meconium aspiration syndrome was most complication which was 67 (67%). Asphyxia found in 55% neonates, 49 (49%) neonates had respiratory distress syndrome, neonatal

jaundice was found in 8 (8%) and sepsis was found in 7 (7%) neonates (Table 4).

**Table No.1: Demographic information of the neonatal**

Variable	No.	%
<b>Gender</b>		
Male	65	65.0
Female	35	35.0
<b>Birth trauma</b>		
Yes	5	5.0
No	95	95.0
<b>NICU admission</b>		
Yes	64	64.0
No	36	36.0
<b>Hospital stay (days)</b>		
<3	51	51.0
4-5	37	37.0
6-7	12	12.0

**Table No.2: Patients biophysical profile (n=100)**

Biophysical Profile Score	No.	%
6/10	10	10.0
8/10	50	50.0
10/10	40	40.0

**Table No.3: Frequency of delivery mode (n=100)**

Delivery Mode	No.	%
Vaginal instrumental	12	12.0
Vaginal spontaneous	20	20.0
C-Section elective	3	3.0
C-Section emergency	65	65.0

**Table No.4: Frequency of neonatal complications**

Complication	No.	%
Sepsis	7	7.0
Neonatal jaundice	8	8.0
Respiratory distress syndrome	49	49.0
Asphyxia	55	55.0
Meconium aspiration syndrome	67	67.0

## DISCUSSION

Inaccurate pregnancy dating is the most common reason to diagnose post-term pregnancy. To assess the gestational age in pregnancy last menstrual period with menstrual series is best physiological landmark. However, there are only few females which are sure about their dates and frequently causes anxiety when they came with postdate.<sup>2</sup> The cause of prolonged pregnancy is unknown. Mostly in obese, nulliparous the post-term pregnancy happens as well as in that females who had post-term pregnancy previously.<sup>4-14</sup> There is high risk of perinatal mortality & morbidity involved in post-term pregnancies which also includes oligohydramnios, sepsis neonatorum, neonatal jaundice meconium aspiration syndrome, fetal distress, fetal birth injury, macrosomia and increase rate of C-

section.<sup>15</sup> Determination of fetal outcome among prolonged pregnancies is the view of this conducted study. In this study, mostly patients (90%) fall under the 20-years to 30-years of age group, which is similar to another study conducted by Oakland.<sup>16</sup> In the study conducted by Oakland showed that 8.60% patients were below the age of thirty four years. According to these findings, in age group of 20-years to 25-years prolonged pregnancy is a common incidence. Although, nulliparous patients are more common for prolonged pregnancies, mostly patients (60%) in our study were multigravida which is similar to the study by Cucco et al.<sup>17</sup> In multigravida patients one of the main recognize factor of prolonged pregnancy is past history of this type of event. Patients who had past history of prolonged pregnancy are at risk of post-term pregnancy in later pregnancy.<sup>18</sup> In our study 40% patients had previous history of post-term pregnancy. In accordance with these findings, special care given by the obstetricians to avoid from later post-term pregnancy and patients may also take special care for prevention from post-term pregnancy.

Although abnormal fetal heart rate reported by many studies, 18% of our patients had deceleration. Prolonged pregnancy is not allied with breech presentation<sup>19</sup> and breech presentation itself allied with increase rate of C-section. In post-term pregnancy caesarean section rate varies quite high as reported incidence varies from 15% to 80%.<sup>20</sup> Nearly 1/3<sup>rd</sup> patients of our study were delivered vaginally, consequently preventing the operation risk. There was caesarean delivery in 68% patients in our study. This high rate is due to that patients were un-booked and directly presented to labour room therefore had no follow-up. Unfavorable cervical findings mostly in our patients at the time of presentation which also contribute to high caesarean section rate in our patients. Prolonged pregnancy is the main reason to increase the fetal morbidity. Due to reduced liquor volume and poor placental reserves, asphyxia and fetal distress is more common.<sup>21</sup> Physiological passage of meconium which occur due to parasympathetic system maturation by forty two weeks of gestation increases the incidence of meconium aspiration.<sup>22</sup> Neonatal complications incidence was quite high in our study e.g. asphyxia, respiratory distress syndrome, meconium aspiration syndrome, sepsis neonatorum and jaundice which are similar incidence of these complications reported internationally.<sup>7-8</sup>

In the absence of other complications, prolonged pregnancy management is controversial. Among obstetricians, there is debate on expectant management of post-term patients versus elective induction of labour.<sup>20,23</sup> Stripping or sweeping the membranes and unprotected coitus may prevent from prolonged pregnancy.<sup>24</sup> To avoid from fetal morbidity close monitoring is required in any case. Careful counseling

concerning the risk and benefits of each component of care should require individual patient management.<sup>25</sup>

## CONCLUSION

Prolonged pregnancy lies high rate of fetal mortality and morbidity with severe complications for mothers. It is concluded from this study that pregnancies should not allow to go post-term as they are associated with higher neonatal mortality and morbidity. Before 40 weeks of gestation induction of labour should be offered to females to prevent from adverse neonatal consequences.

### Author's Contribution:

Concept & Design of Study: Ashba Anwer  
 Drafting: Nazia Tufail  
 Data Analysis: Asma Mudassir  
 Revisiting Critically: Ashba Anwer  
 Final Approval of version: Ashba Anwer

**Conflict of Interest:** The study has no conflict of interest to declare by any author.

## REFERENCES

1. American College of Obstetricians and Gynecologists. Practice bulletin no. 146: Management of late-term and postterm pregnancies. *Obstet Gynecol* 124(2pt1): 390-6.
2. Norwitz ER, Snegovskikh W, Caughey AB. Prolonged pregnancy. *Clin Obstet Gynecol* 2007; 50(2):547-57.
3. Bennett KA, Crane JMG, O'Shea P, Lacelle MD, Hutchens D, Copel J. First trimester ultrasound screening is effective in reducing post-term induction rates: a randomized controlled trial. *Am J Obstet Gynecol* 2004; 190:1077-81.
4. Kistka ZA, Palomar L, Lee KA, Boslaugh SE, Wangler MF, Cole FS, et al. Radical disparity in the frequency of recurrence of preterm birth. *Am J Obstet Gynecol* 2007; 196:131.e1-6.
5. Olesen AW, Westergaard JG, Olsen J; Prenatal risk indicators of a prolonged pregnancy. The Danish Birth Cohort 1998-2001. *Acta Obstet Gynecol Scand* 2006; 85(11):1338-41.
6. Gardosi J, Vanner T, Francis A. Gestational age and induction of labour for prolonged pregnancy. *Br J Obstet Gynaecol* 1997; 104(7):792-7.
7. Savitz D, Terry JW, Dole N, Thorp JM, Siega-Riz AM, et al. Comparison of pregnancy dating by last menstrual period, ultrasound scanning, and their combination. *Am J Obstet Gynecol* 2002; 187: 1660-6.
8. Smith GCS. Life-table analysis of the risk of perinatal death at term and post term in singleton pregnancies. *Am J Obstet Gynecol* 2001; 184: 489-96.

9. Olesen AW, Westergaard JG, Olsen J. Perinatal and maternal complications related to postterm delivery: a national register-based study, 1978–1993. *Am J Obstet Gynecol* 2003;189:222–7.
10. Hilder L, Costeloe K, Thilaganathan B. Prolonged pregnancy: evaluating gestation-specific risks of fetal and infant mortality. *Br J Obstet Gynaecol* 1998;105:169–73.
11. Royal College of Obstetricians and Gynecologists. Induction of labour. Evidence based clinical Guideline No. 9. London: RCOG Press, 2009.
12. Dasari P, Niveditta G, Raghavan S. The maximal vertical pocket and amniotic fluid index in predicting fetal distress in prolonged pregnancy. *Int J Gynaecol Obstet* 2007;96(2):89-93.
13. Treger M, Hallak M, Silberstein T, Friger M, Katz M, Mazor M. Post-term pregnancy: should induction of labor be considered before 42 weeks? *J Matern Fetal Med* 2002;11:50–3.
14. Gelisen O, Caliskan E, Dilbaz S, Ozdas E, Dilbaz B, Ozdas E, et al. Induction of labor with three different techniques at 41 weeks of gestation or spontaneous follow-up until 42 weeks in women with definitely unfavorable cervical scores. *Eur J Obstet Gynecol Reprod Biol* 2005;120(2):164–9.
15. Roach VJ, Rogers MS. Pregnancy outcome beyond 41 weeks gestation. *Int J Gynecol Obstet* 1997; 59:19–24.
16. Cleary-Goldman J, Bettles B, Robinson JN, Norwitz E, D'Alton ME, Schulkin J. Postterm pregnancy: practice patterns of contemporary obstetricians and gynecologists. *Am J Perinatol* 2006;23(1):15-20.
17. Aaron B, Caughey MD. Maternal and obstetric complications of pregnancy are associated with increasing gestational age at term. *Am J Obstet Gynecol* 2009;200:683 -5.
18. Cucco C, Osbrone MA, Cibils LA. Maternal-fetal outcomes in prolonged pregnancy. *Am J Obstet Gynecol* 1989; 161:916-20.
19. James C, George SS, Guanekar N, Seshadri L. Management of prolonged pregnancy: arandomized trial of induction and antepartum foetal monitoring. *Nat J Ind* 2001;14:270–3.
20. Hannah ME. Postterm pregnancy: should all women have labour induced? A review of the literature. *Fetal Maternal Med Rev* 1993;5:3–17.
21. Heimstad R, Skogvoll E, Mattson L, Johansen OJ, Eik-Nes SH, Salvesen KA. Induction of labor or serial antenatal fetal monitoring in post-term pregnancy. *Obstet Gynecol* 2007;109:609–17.
22. Caughey AB, SnegovskikhVV, Norwitz ER. Postterm pregnancy: how can we improve outcomes? *Obstet Gynecol Surv* 2008; 63(11): 715-24.
23. Mogen I, Stenlund H, Hodberg U. Recurrence of prolonged pregnancy. *Int J Epidemiol* 1999; 28(2):253-7.
24. Herabutya Y, Prasertsawat PO, Tongyai T, Isarangura Na Ayudhya N. Prolonged pregnancy: the management dilemma. *Int J Gynecol Obstet* 1992;37:253–8.
25. de Miranda E, van der Bom JG, BonselGJ, Bleker OP, Rosendaal FR. Membrane sweeping and prevention of post-term pregnancy in low-risk pregnancies: a randomized controlled trial. *BJOG* 2006;113:402–8.