

Is Ultrasound Detect Renal Infections?

Detection of
Renal Infection
on Ultrasound

Nisar Ahmed Shaikh¹, Hameed-ur-Rahman Bozdar³, Akber Ali Soomro² and Malik Hussain Jalbani¹

ABSTRACT

Objective: To assess culture and sensitivity of renal infection patients detected on ultrasound

Study Design: Prospective cohort study

Place and Duration of Study: This study was conducted at the Rimsha Medical Center Dadu from January 2012 to June 2015.

Materials and Methods: Hundred patients suffering for renal infection detected on ultrasound were included in the study. Diabetic and urolithiasis patients excluded from the study. Urine culture and sensitivity of renal infection patients detected on ultrasound carried from collection point of diagnostic and research laboratory Liaquat University of Medical & Health Sciences Jamshoro/Hyderabad in Dadu.

Results: Out of 100 patients 65 (65%) were female and 35(35%) male. The age of the patients ranged from 15 to 45 years with mean age 28.9 SD±8.8 The presenting features were lumbar pain with on and off fever in 60(60%) cases, dysuria in 15(15%), and in 35(35%) cases was asymptomatic. The renal infection detected in ultrasound bilaterally in 30(30%), Right sided in 35(35%), left sided in 25(25%) and chronic pyelonephritis (change of small size kidney with increase echogenicity and small size kidney with irregular border) in 10(10%) cases. Culture and sensitivity seen positive in 27(27%) cases. Pyuria and haematuria seen in 17(17%). The commonest micro-organism detected E.coli in 60(60%), Klebsiella 15(15%), Proteus 5(5%), Enterococcus 5(5%), Staphylococcus saprophyticus 5(5%), Streptococcus 7(7%), Pseudomonas 3(3%) and insignificant mixed bacterial growth seen in 7 cases. Drug sensitivity seen in 100% with meronam, piperacillin/tazobactam, aztreonam. Nitrofurantoin and amikacin 90% Amoxiclavulanic acid in fusidic acid, ceftriaxone, cotrimoxazole, ofloxacin, cefuroxime, cefixime 75% and resistance pattern seen more in ceftazidime, gentamycin, ciprofloxacin and ampicillin.

Conclusion: Renal infection detected on ultrasound even with insignificant pyuria and haematuria should not be neglected because having significant positive culture and sensitivity report findings.

Key Words: Renal infection, ultrasound, culture and sensitivity

Citation of article: Shaikh NA, Bozdar HR, Soomro AA, Jalbani MH. Is Ultrasound Detect Renal Infections? Med Forum 2016;27(10):16-18.

INTRODUCTION

Renal infection (bacterial pyelonephritis) refers to any inflammation affecting the renal interstitium. Patients most often affected are females from 15- to 40-years-old. Predisposing conditions include neuropathic bladder, prolonged catheter drainage, urine reflux, bladder malignancy, urinary bladder outlet obstruction, clinical Benign prostatic enlargement, calculus disease, altered host resistance, congenital anomalies, analgesic abuse, diabetes, sexual activity and pregnancy.^{1,8} Renal infection is presenting fairly specific and classical in most cases, consisting of rapid onset of high fevers and flank pain and tenderness.

In many instances less specific or non-urinary symptoms and signs may also be present which may lead to clinical confusion.^{2,4} A renal infection requires prompt medical attention. If not treated properly, a renal infection can permanently damage kidneys by renal abscess, renal infarction, necrosis, scarring, chronic renal impairment and hypertension. The most commonly implicated micro organisms are E. coli (most common), Klebsiella, Proteus, Enterobacter and Pseudomonas.^{8,14}

Ultrasound is insensitive to the changes of acute pyelonephritis, with most patients having normal scan, and abnormalities only identified in 25% of cases. Possible features like particulate matter in the collecting system, gas bubbles (emphysematous pyelonephritis), abnormal echogenicity of the renal parenchyma and focal/segmental hypoechoic regions, increase diameter > 50 millimeter in transverse section with prominent pyramids seen in kidney. However ultrasound also useful in assessing for local complications such as hydronephrosis, renal abscess formation, renal infarction, perinephric collections, and thus guiding

¹. Department of Urology / Pathology², CMC, Larkana, Sindh

³. Department of Urology, KMC&H, Khairpur, Sindh

Correspondence: Nisar Ahmed Shaikh, Asstt. Prof. of Urology, CMC, Larkana, Sindh

Contact No: 0300-3428026

Email: dnmisarshaikh@yahoo.com

Received: July 27, 2016;

Accepted: August 30, 2016

management.^{1,7} Still some radiologist use term renal infection when above finding is noted in ultrasound but still that's debatable. So our aim of study to asses culture and sensitivity of renal infection patients detected on ultrasound

MATERIALS AND METHODS

A Prospective cohort study of 100 cases of renal infection carried out at Rimsha medical center Dadu from Jan. 2012 to June 2015. The study was approved by the ethical review committee and informed consent was personally obtained from the patients. Hundred patients suffering for renal infection detected on ultrasound were included in the study. Neuropathic bladder, prolonged catheter drainage, urine reflux, bladder malignancy, urinary bladder outlet obstruction, clinical Benign prostatic enlargement and calculus disease patients excluded from the study. Urine culture and sensitivity of renal infection patients detected on ultrasound carried from collection point of diagnostic and research laboratory liaqut university of medical & health sciences Jamshoro/Hyderabad in Dadu, in all cases to asses pattern of urine culture and sensitivity.

RESULTS

Out of 100 patients, 75(75%) were female and 25(25%) male. The age of the patients ranged from 15 to 45 year with mean age 28.9SD=8.8 The presenting features were lumbar pain with on & off fever in 60(60%) cases, dysuria in 15(15%), and in 35(35%) cases were asymptomatic. The acute renal infection /acute pyelonephritis suspected from increased echotexture of kidney, increase diameter > 50 millimeter in transverse section with prominent pyramids seen in bilaterally in 30(30%), Right sided in 35(35%), left sided in 25(25%) and chronic pyelonephritis in 10(10%) cases. culture and sensitivity seen positive in 27(27%) cases. Pyuria and haematuria seen in 17(17%). Urine culture and sensitivity showed commonest micro-organism detected E.coli in 60(60%), Klebsiella 15(15%), Proteus 5(5%), Enterococcus 5(5%), staphylococcus saprophyticus 5(5%), streptococcus 7(7%), Pseudomonas 3(3%) and insignificant mixed bacterial growth seen in 7(7%). Drug sensitivity seen in 100% with meronam, piperacillin/tazobactam. Nitrofurantoin and amikacin in 90%. Amoxi- clavulanic acid and fusidic acid, ceftriaxone, cotrimoxazole, ofloxacin, cefixime 75% and resistance pattern seen more in ciprofloxacin, penicillin and ampicillin.



Figure No.1. Various ultrasounds show renal infection

DISCUSSION

The aim of imaging in renal infection is to detect conditions that must be treated in order to avoid immediate deterioration or recurrences, and long-term kidney damage.^{4,9} In our study of 100 patients of renal infection suspected on ultrasonography 75(75%) were female and 25(25%) male. The age of the patients ranged from 15 to 45 years with mean age 28.9 SD±8.8 which is comparable to Bjerklund TE study.⁷ The presenting features were lumbar pain with on and off fever in 60(60%) cases, dysuria in 15(15%), and in 35(35%) cases was asymptomatic. The renal infection seen in bilaterally in 30(30%), right sided 35(35%), left sided 25(25%) and chronic pyelonephritis increased echogenicity, small size kidney with irregular border) 10(10%) cases. Pyuria and haematuria seen in 17(17%) which is also comparable to secondary data.^{5,10} Our study hypothesis was that there is co-relation of renal infection detected on ultrasound with UTI, is some

thing true but verbal communication with various radiologists are in favour of null hypothesis that there is no co-relation among renal infection detected on ultrasound with UTI so we collect 100 sample of urine for culture and sensitivity of renal infection detected by ultrasound and found culture and sensitivity positive in 27(27%) cases with P-value 0.02 which is statistically significant hence adequate evidence against the null hypothesis. The commonest micro-organism detected E.coli in 60(60%), Klebsiella 15(15%), Proteus 5(5%), Enterococcus 5(5%), staphylococcus saprophyticus 5(5%), streptococcus 7(7%), Pseudomonas 3(3%) and insignificant mixed bacterial growth seen in 7(7%). Drug sensitivity seen in 100% with meronam, tazobactam. Nitrofurantoin and amikacin seen in 90%. Cavulanic acid and fusidic acid, ceftriaxone, cotrimoxole, ofloxin, cefixime seen in 75% and resistance pattern seen more in ciprofloxacin, penicillin and ampicillin which is comparable to Rossleigh MA study.¹¹⁻¹³ So we recommended that renal infection

detected on ultrasound even with insignificant pyuria and haematuria should not be neglected because our study showed 27(27%) positive cases of culture and sensitivity cases with p-value 0.02 which is statistically significant. However further studies needed to confirm that renal infection detected on ultrasound have correlation with positive cases of culture and sensitivity.

CONCLUSION

Renal infection detected on ultrasound even with insignificant pyuria and haematuria should not be neglected because having significant positive culture and sensitivity report findings.

Conflict of Interest: The study has no conflict of interest to declare by any author.

REFERENCES

1. Kawashima A, LeRoy AJ. Radiologic evaluation of patients with renal infections. *Infect Dis Clin North Am* 2003;17(2):433-56.
2. Naber B, Bergman MC, Bisho TE, Bjerklund J, Botto B, Lobel, et al. EAU guidelines for the management of urinary and male genital tract infections. *Eur Urol* 2001;40:576-588.
3. Cergole MC, Pignatari AC and Guth BE.. Adhesion, biofilm and genotypic characteristics of antimicrobial resistant *Escherichia coli* isolates. *Braz J Microbiol* 2015;46(1):167-71.
4. Craig BJ, Wagner MD and Travis. Pyelonephritis: radiologic-pathologic review *Int J Infect Dis* 2013;17(5):254-85.
5. Narchi H and Donovan R. Renal power Doppler ultrasound does not predict renal scarring after urinary tract infection. *Scott Med J* 2008; 53(4):7-10.
6. Rossleigh MA. Renal infection and vesico-ureteric reflux. *Semin Nucl Med* 2007;37(4):261-8.
7. Bjerklund TE. Diagnosis and imaging in urinary tract infections. *Curr Opin Urol* 2002;12(1):39-43.
8. Hellerstein S. Acute urinary tract infection, evaluation and treatment. *Curr Opin Pediatr* 2006; 18(2):134-8.
9. Nelson CP, Johnson EK, Logvinenko T, Chow JS. Ultrasound as a screening test for genitourinary anomalies in children with UTI. *Pediatrics* 2014; 133(3):394-403.
10. Riccabona M. Imaging in childhood urinary tract infection. *Radiol Med* 2016;121(5):391-401.
11. Kurtz MP, Chow JS, Johnson EK, Rosoklija I, Logvinenko T, Nelson CP. Imaging after urinary tract infection in older children and adolescents. *J Urol* 2015;193(5 Suppl):1778-82.
12. Guedj R, Escoda S, Blakime P, Patteau G, Brunelle F, Cheron G. The accuracy of renal point of care ultrasound to detect hydronephrosis in children with a urinary tract infection. *Eur J Emerg Med* 2015;22(2):135
13. Johnson CE, DeBaz BP, Shurin PA, DeBartolomeo R. Renal ultrasound evaluation of urinary tract infections in children. *Pediatrics* 1986;78(5):871-8.
14. Wallace SS, Zhang W, Mahmood NF, Williams JL, Cruz AT, Macias CG, et al. Renal Ultrasound for Infants Younger Than 2 Months With a Febrile Urinary Tract Infection. *AJR Am J Roentgenol* 2015;205(4):894-8.