

Screening of Diabetes and HIV Infection in Newly Diagnosed Pulmonary Tuberculosis Patients

Rizwan Hafeez, Sajjad Ahmed Khan, Azzan Mujahid and Abdullah Irshad

ABSTRACT

Objective: Aim of the study was to investigate the prevalence of diabetes and HIV infection by screening in newly diagnosed pulmonary tuberculosis patients.

Study Design: Cross sectional study.

Place and Duration of Study: This study was conducted at the Pulmonology Department of Shahida Islam Medical College, Lodhran from 10 June 2016 to 10 June 2017.

Materials and Methods: This study was conducted after taking approval from the institutional ethical committee. Total 159 newly diagnosed pulmonary tuberculosis patients were recruited by non-probability consecutive sampling. Mean and SD was calculated for numerical variables like age, BMI, fasting blood glucose (FBG), Hemoglobin and monthly income. Frequency and percentages were calculated for categorical variables like gender, smoking status, educational status, alcohol consumption, prevalence of HIV and diabetes, Chi square test was used to check significant difference in PTB and Non PTB patients. P value of <0.05 was taken as significant.

Results: Significant difference was found between age ($p=0.000$), BMI ($p=0.000$), FBG ($p=0.000$), hemoglobin ($p=0.000$), gender ($p=0.002$) and alcohol consumption ($p=0.034$), in groups, except education status ($p=0.090$) and monthly income ($p=0.068$). The main outcome variables of this study were HIV infection and diabetes. HIV infection was noted in 3.1% ($n=5$) and 1.9% ($n=3$) patients for PTB and non-PTB group respectively. While, 8.2% ($n=13$) and 4.4% ($n=4$) patients suffered from diabetes, for both the groups respectively.

Conclusion: The results of our study revealed that on screening of newly diagnosed pulmonary tuberculosis patients with HIV infection and diabetes, their prevalence was high as compared to non-pulmonary tuberculosis group. But this difference was statistically non significant (P value 0.474 and 0.166).

Key Words: Pulmonary tuberculosis, Diabetes, HIV infection.

Citation of articles: Hafeez R, Khan SA, Mujahid A, Irshad A. Screening of Diabetes and HIV Infection in Newly Diagnosed Pulmonary Tuberculosis Patients. *Med Forum* 2018;29(1):51-55.

INTRODUCTION

Diabetes is metabolic disorder. Insulin resistance is the main pathogenesis in development of type 2 diabetes mellitus. Type 2 diabetes usually affects elderly patients. Type 2 diabetes mellitus is accelerating pandemic throughout the world¹. 230 million population of the world suffer from diabetes. It has been estimated that this population will reach up to 552 million until 2030². It is recorded that 471 billion USD were spent to treat diabetes in 2012³. For the developing countries, it is a great socio economic burden. It is alarming that prevalence of diabetes has significantly risen in past two decades. In 2011 diabetes prevalence was ranging from 7.6% (5.2 million people) to 11% in Pakistan and it is expected to reach up to 15% (14 million) until 2030⁴.

Department of Pulmonology, Bahawal Victoria Hospital, Bahawalpur.

Correspondence: Dr. Sajjad Ahmed Khan, Ex-House Officer, Department of Pulmonology, Bahawal Victoria Hospital, Bahawalpur.

Contact No: 0303-7200923

Email: dr.sajjadahmed084@gmail.com

Received: August, 2017;

Accepted: November, 2017

Pakistan is on number 7 place in list of countries having diabetes mellitus. It is estimated that if this trend continues than it will become number 4. Overall ratio of diabetes is that its prevalence in urban population is 22.04% and in rural area is 17.15% in Pakistan⁵. Pakistan health care system is overburdened and under resource in dealing with this situation. Considering the present situation, it demands a national effort in not only treating but more on preventing the diabetes mellitus. Many complications are associated with diabetes mellitus. Cardiovascular diseases due to diabetes is main cause of premature illness and deaths⁶. It is main cause of renal failure and blindness. It is associated with amputations due diabetic foot development.

Tuberculosis is one of the main public health problem worldwide⁷. It has been investigated that one third population of world is suffering from mycobacterium infection⁸. Patients of pulmonary tuberculosis clinically present mainly with cough, fever, hemoptysis and weight loss. Diabetes is well known risk factor for tuberculosis⁹. World health organization has classified that eight of ten countries which have highest burden of diabetes also have high burden of tuberculosis¹⁰. In developing countries such as Pakistan, India,

Bangladesh and Brazil where tuberculosis is endemic, the burden of these two diseases and interaction between them will be more dangerous. World health organization has declared global epidemic both diabetes and tuberculosis¹¹.

Relationship and association of HIV with tuberculosis is great socio economic threat and public health problem in developing countries. World health organization had estimated in 1992 that about four million population of the world had been infected by both HIV and mycobacterium tuberculosis¹². Association between HIV and tuberculosis is evident by high incidence of tuberculosis in HIV infected patients. In developing countries where health care systems are already overburdened these two epidemics impose grave social and medical implications.

To best of our knowledge in south Punjab of Pakistan no study had been conducted to investigate the prevalence of diabetes and HIV infection in newly diagnosed pulmonary tuberculosis patients. So this study was conducted to investigate the prevalence of these diseases in newly diagnosed pulmonary tuberculosis patients. It will provide a base for further research on this issue and will create awareness. Study done by Qiuzhen Wang et al. was taken as reference study¹³.

MATERIALS AND METHODS

This cross sectional study was conducted in the Pulmonology department of Shahida Islam Medical College, Lodhran. From 10 June 2016 to 10 June 2017 after taking approval from the institutional ethical committee. Total 159 newly diagnosed pulmonary tuberculosis patients were recruited by non-probability consecutive sampling. To study the prevalence of diabetes and HIV infection in non-tuberculosis group, by cluster random sampling 159 subjects were recruited from same communities as that of tuberculosis patients. Written permission of study was signed by each individual recruited in study. Exclusion criteria of study were: 1) patients with age < 18 years of age, 2) patients who had already taken anti tuberculosis drugs, 3) patients who were known to have diabetes and HIV infection 4) patients who had any history of oncological disorder and 5) patients who had other immunosuppressive conditions. Sample size of study was calculated by using a reference study done by Qiuzhen Wang et al. for which confidence interval was taken as 95%, study strength and odd ratio of diabetes in newly diagnosed pulmonary tuberculosis patients was 3.17%¹³.

All patients were recruited from outpatient department. Detailed medical history of individual was taken and through clinical examination was conducted to record any complication of the disease and to assess any sign of other immunosuppressive disorder. Blood pressure and vitals were recorded at time of pulmonary

tuberculosis patients. Blood samples were also taken to check glucose levels and hemoglobin level. Personal information like age, gender, living area, income, body mass index duration of symptoms was taken by filling the Performa.

Pulmonary tuberculosis was diagnosed according to the guidelines of National Tuberculosis Program. Patients were investigated with smear examination who were suspected to have pulmonary tuberculosis. Patients who were smear negative but having clinical and radiological suspicion were also labelled as pulmonary tuberculosis after discussing it with radiologist and chest consultants.

Diabetes was diagnosed by using standard method of checking fasting plasma glucose levels as recommended by World Health Organization. Venous blood was used after overnight fasting to check plasma glucose level. Patients having plasma glucose level greater than 126mg/dl were diagnosed as diabetics. Screening of HIV infection was done in each individual recruited in study by using ELISA method.

Data was analyzed by using SPSS volume 23. Quantitative variables like age, body mass index, plasma glucose levels and age were statistically analyzed by their mean and standard deviation and t-test was applied to check the significance. Qualitative variables like gender, living area, HIV infection, diabetes and income were statistically analyzed by frequency and percentage and chi-square test was used to check the significance. P value < 0.05 was considered as statistically significant.

RESULTS

In this study, a total number of 100% (n=318) patients were included, divided into two equal groups, 50% (n=159) in each i.e. PTB and non-PTB group. The Mean±S.D of age, BMI, FBG and hemoglobin of the patients of PTB group was 46.14±2.92 years, 21.16±2.46 kg/m², 6.02±1.15 and 10.82±1.74 g/dl respectively. While, the Mean±S.D of age, BMI, FBG and hemoglobin of the patients of non-PTB group was 48.48±3.20 years, 23.31±2.24 kg/m², 5.13±1.72 and 12.53±1.64 g/dl respectively. Gender distribution, in PTB group, was observed as 74.8% (n=119) males and 25.2% (n=40) females. While, in non-PTB group, there were 58.5% (n=93) males and 41.5% (n=66) females. There were 28.3% (n=45) and 13.2% (n=21) smokers in PTB and non-PTB group respectively. Education status of PTB group was observed as 35.2% (n=56) illiterate, 39% (n=62) matric and 25.8% (n=41) graduate or above. However, in non-PTB group, 37.1% (n=59) illiterate, 28.3% (n=45) matric and 34.6% (n=55) graduate or above. Income level of the patients of PTB group observed as 35.2% (n=56) patients had <200000 rupees/year and 64.8% (n=103) had >200000 rupees/year. While, 25.8% (n=41) patients of non-PTB group had <200000 rupee/year income and 74.2%

(n=118) had >200000 rupee/year income. There were 10.7% (n=17) and 4.4% (n=7) patients used alcohol of PTB and non-PTB group respectively. Significant difference was found between age (p=0.000), BMI (p=0.000), FBG (p=0.000), hemoglobin (p=0.000), gender (p=0.002) and alcohol consumption (p=0.034), in groups, except education status (p=0.090) and monthly income (p=0.068). (Table. 1& 3).

The main outcome variables of this study were HIV infection and diabetes. HIV infection was noted in 3.1% (n=5) and 1.9% (n=3) patients for PTB and non-PTB group respectively. While, 8.2% (n=13) and 4.4% (n=4) patients suffered from diabetes, for both the groups respectively. No significant difference was found between HIV (p=0.474) and diabetes (p=0.166), in groups. (Table. 2). (Figure. 1).

Table No. 1: Characteristics of PTB and Non-PTB patients

Variable	PTB (n=159)	Non-PTB (n=159)	Test of Sig.
Age	46.14± 2.92 years	48.48± 3.20 years	t=-6.81, p=0.000
BMI	21.16± 2.46 kg/m ²	23.31± 2.24 kg/m ²	t=-8.13, p=0.000
Fasting blood glucose	6.02± 1.15	5.13± 1.72	t=-8.94, p=0.000
Hemoglobin	10.82± 1.74 g/dl	12.53± 1.64 g/dl	t=5.39, p=0.000
Gender	M=74.8%, F=25.2%	M=58.5%, F=41.5%	$\chi^2=9.56$, p=0.002
Smoking Status	Smokers=2 8.3%	Smokers=13 13.2%	$\chi^2=11.01$, p=0.001
Education Status	Illiterate=3 5.2% Matric=39% graduate or above=25.8%	Illiterate=37 37.1% Matric=28.3% graduate or above=34.6%	$\chi^2=4.82$, p=0.090
Monthly Income	<200000=3 5.2% >200000=6 4.8%	<200000=25 8% >200000=74 74.2%	$\chi^2=3.33$, p=0.068
Alcohol Consumption	10.7%	4.4%	$\chi^2=4.51$, p=0.034

Table No. 2: Distribution of HIV and Diabetic in groups

Variable	PTB (n=159)	Non-PTB (n=159)	Test of Sig.
HIV	3.1%	1.9%	$\chi^2=0.513$,p=0.474
Diabetic	8.2%	4.4%	$\chi^2=1.92$,p=0.166




Figure No.1: Distribution of HIV and Diabetic in groups

Table No. 3: Association of groups between effect modifiers

Variable	Groups		Total	Chi-Square P-value
	PTB	Non-PTB		
Gender	Male	119	93	0.002
	Female	40	66	
Smoking	Yes	45	21	0.001
	No	114	138	
Educational Status	Illiterate	56	59	0.000
	Matric	62	45	
	Graduate or above	41	55	
Income	<200000	56	41	0.068
	>200000	103	118	
Alcohol	Yes	17	7	0.034
	No	142	152	
Diabetic	Yes	13	7	0.166
	No	146	152	
HIV	Yes	5	3	0.474
	No	154	156	

DISCUSSION

Results of study showed that prevalence of diabetes mellitus on screening in newly diagnosed pulmonary tuberculosis patients was higher (8.2%) than in non-pulmonary tuberculosis (4.4%) group. Similarly study also showed that HIV infection on screening was higher (3.1%) in newly diagnosed pulmonary tuberculosis patients than in non-pulmonary tuberculosis (1.9%) group. But this difference was statistically non-significant with P value 0.474 and 0.166.

To best of our knowledge there was no data available about prevalence of diabetes and HIV infection among newly diagnosed pulmonary tuberculosis patients as well as comparative data from non-pulmonary tuberculosis controls from the same community collected at same time. The study was designed to discover the hidden prevalence of HIV infection and diabetes mellitus in newly diagnosed pulmonary tuberculosis patients in Pakistan generally and in our local geographical area specially. Strength of this study is that, to define diabetes mellitus accurately, primary data obtained from newly diagnosed

pulmonary tuberculosis patients and non-pulmonary tuberculosis group was used. While in other studies, either the prevalence of diabetes mellitus in non-pulmonary tuberculosis patients was not reported or it was estimated from secondary data with inherent biases¹⁴⁻¹⁵. Study done by Tahir et al. showed that prevalence of diabetes mellitus among pulmonary tuberculosis patients is 14.8%¹⁶. Study done by Jabbaret al. in Karachi and found that tuberculosis was ten times more prevalent in diabetic patients than in non-diabetic patients and prevalence was also increased with increase in duration of diabetes mellitus¹⁷. Another study carried out by Zheng et al. found that prevalence of diabetes mellitus in tuberculosis was ranging from 5% to more than 50%¹⁸. Results of study done by Mukhtar et al. showed that diabetes is associated with unfavorable outcomes in pulmonary tuberculosis patients¹⁹. Study of Alavi et al. concluded that pulmonary tuberculosis should be considered at top of the list in every diabetic patients who presented with cough, sputum and dyspnea with cavitation and miliary mottling on chest x ray²⁰. Many studies had been done to evaluate the prevalence and association of tuberculosis in already known HIV infected patients. But data is very limited about screening of HIV infection in newly diagnosed pulmonary tuberculosis patients. It has been noted that incidence of HIV infection is growing in our region. Study done by Giri et al. showed that 17% HIV infected patients had tuberculosis co-infection²¹. Study carried out by Chisti et al. found that pulmonary tuberculosis can be a common cause of acute pneumonia in severely malnourished or HIV infected children under five years old²².

CONCLUSION

The results of our study revealed that on screening of newly diagnosed pulmonary tuberculosis patients with HIV infection and diabetes, their prevalence was high as compared to non-pulmonary tuberculosis group. But this difference was statistically non significant (P value 0.474 and 0.166).

Author's Contribution:

Concept & Design of Study:	Rizwan Hafeez
Drafting:	Sajjad Ahmed Khan
Data Analysis:	Azzan Mujahid, Abdullah Irshad
Revisiting Critically:	Rizwan Hafeez, Sajjad Ahmed Khan
Final Approval of version:	Rizwan Hafeez

Conflict of Interest: The study has no conflict of interest to declare by any author.

REFERENCES

1. Mahishale V, Avuthu S, Patil B, Lolly M, Eti A, Khan S. Effect of Poor Glycemic Control in Newly Diagnosed Patients with Smear-Positive Pulmonary Tuberculosis and Type-2 Diabetes Mellitus. *Iran J Med Sci* 2017;42(2):144.
2. Guo SH, Chang HK, Lin CY. Impact of Mobile Diabetes Self-Care System on patients' knowledge, behavior and efficacy. *Computers in Indust* 2015;69:22-9.
3. He L, Meng S, Germain-Lee EL, Radovick S, Wondisford FE. Potential biomarker of metformin action. *J Endocrino* 2014;221(3):363-9.
4. Hussain A, Ali I. Diabetes mellitus in Pakistan: A major public health concern. *Arch Pharmacy Pract* 2016;7(1):30.
5. Bahadar H, Mostafalou S, Abdollahi M. Growing burden of diabetes in Pakistan and the possible role of arsenic and pesticides. *J Diabet Metabolic Dis* 2014;13(1):117.
6. Li G, Zhang P, Wang J, An Y, Gong Q, Gregg EW, et al. Cardiovascular mortality, all-cause mortality, and diabetes incidence after lifestyle intervention for people with impaired glucose tolerance in the Da Qing Diabetes Prevention Study: a 23-year follow-up study. *Lancet Diabetes Endocrinol* 2014; 2(6):474-80.
7. Eurosurveillance Editorial Team. WHO publishes Global tuberculosis report 2013. *Eurosurveillance* 2013;18(43):20615.
8. Gopal R, Monin L, Slight S, Uche U, Blanchard E, Junecko BA, et al. Unexpected role for IL-17 in protective immunity against hypervirulent *Mycobacterium tuberculosis* HN878 infection. *PLoS Pathogens* 2014;10(5):e1004099.
9. Root HF. The association of diabetes and tuberculosis. *N Eng J Med* 1934;210(3):127-47.
10. Lönnroth K, Roglic G, Harries AD. Improving tuberculosis prevention and care through addressing the global diabetes epidemic: from evidence to policy and practice. *Lancet Diabetes Endocrinol* 2014;2(9):730-9.
11. Kapur A, Harries AD, Lönnroth K, Wilson P, Sulistyowati LS. Diabetes and tuberculosis co-epidemic: the Bali Declaration. *Lancet Diabetes Endocrinol* 2016;4(1):8-10.
12. Narain JP, Raviglione MC, Kochi A. HIV-associated tuberculosis in developing countries: Epidemiology and Strategies for Prevention. *Tuber Lung Dis* 1992;73(6):311-21.
13. Wang Q, Ma A, Han X, Zhao S, Cai J, Ma Y, et al. Prevalence of type 2 diabetes among newly detected pulmonary tuberculosis patients in China: a community based cohort study. *PLoS One* 2013; 8(12):e82660.

14. Restrepo BI, Camerlin AJ, Rahbar MH, Wang W, Restrepo MA, Zarate I, et al. Cross-sectional assessment reveals high diabetes prevalence among newly-diagnosed tuberculosis cases. *Bull World Health Organ* 2011;89(5):352-9.
15. Viswanathan V, Kumpatla S, Aravindalochanan V, Rajan R, Chinnasamy C, Srinivasan R, et al. Prevalence of diabetes and pre-diabetes and associated risk factors among tuberculosis patients in India. *PloS One* 2012;7(7):e41367.
16. Tahir Z, Akhtar AM, Yaqub T, Mushtaq MH, Javed H. Diabetes mellitus among tuberculosis patients: a cross sectional study from Pakistan. *AfrHealth Sci* 2016;16(3):671-6.
17. Jabbar A, Hussain SF, Khan AA. Clinical characteristics of pulmonary tuberculosis in adult Pakistani patients with co-existing diabetes mellitus. *East Mediterr Health J* 2006;12(5):522-7.
18. Zheng C, Hu M, Gao F. Diabetes and pulmonary tuberculosis: a global overview with special focus on the situation in Asian countries with high TB-DM burden. *Glob Health Action* 2017;10(1):1-11.
19. Mukhtar F, Butt ZA. Cohort profile: the diabetes-tuberculosis treatment outcome (DITTO) study in Pakistan. *BMJ Open* 2016;6(12):e012970.
20. Alavi SM, Khoshkho MM, Salmanzadeh S, Eghtesad M. Comparison of epidemiological, clinical, laboratory and radiological features of hospitalized diabetic and non-diabetic patients with pulmonary tuberculosis at razi hospital in ahvaz. *Jundishapur J Micro* 2014; 7(9): e12447.
21. Giri PA, Deshpande JD, Phalke DB. Prevalence of pulmonary tuberculosis among HIV positive patients attending antiretroviral therapy clinic. *N Am J Med Sci* 2013;5(6):367.
22. Chisti MJ, Ahmed T, Pietroni MA, Faruque AS, Ashraf H, Bardhan PK, et al. Pulmonary tuberculosis in severely-malnourished or HIV-infected children with pneumonia: a review. *J Health Popul Nutr* 2013;31(3):308.