

Evaluation of Liver Injuries in DHQ Teaching Hospital Bannu

Ajmal Shah Bukhari¹, Shabir Hussain² and Mir Hamza³

ABSTRACT

Objective: To determine the major injury patterns, outcomes and management options of liver trauma in a tertiary care setup in District Headquarter Teaching Hospital Bannu, Pakistan.

Study Design: Retrospective / clinical study

Place and Duration of Study: This study was conducted at the Department of Surgery, District Headquarter Teaching Hospital Bannu from 1st January 2015 till 31st December 2016.

Materials and Methods: It was carried out consisted of 60 patients with liver trauma, 57 males and 3 female, with the mean age 31.46 years. Data regarding age, sex, mode and type of injuries were taken and analyzed. Inclusion criteria included age group equals or more than 13 years of age with diagnosis of liver trauma, patients penetrating and non-penetrating traumatic injury to liver, patients with blunt and sharp injury to liver. Exclusion criteria included all the patients' less than 13 years of age, patients with pre-existing liver disease i.e. cirrhosis, tumors, hepatitis etc, Patients who have previously undergone hepatic surgery.

Results: The incidence of liver trauma due to non-penetrating injuries was 46(76.6%) while due to penetrating injuries 14 (23.3%), In all cases of blunt injuries, 40 patients (86.9%) were due to road traffic accidents, and 6 patients (13.0%) were due to assaults. 16 patients who were haemodynamically stable, were managed Medically with strict vital monitoring, input/output charting and repeated examinations. 44 patients who were haemodynamically unstable despite aggressive resuscitation and were managed surgically. 9 patients (20%) were treated by simple suture, 15 patients (34%) were treated by suture and perihepatic packing, and 20 patients (45%) were treated by perihepatic packing. Patient who underwent perihepatic packing were re explored after 48 hours. abdomen was washed, drains were put in and abdomen was closed permanently.

Conclusion: Non-penetrating liver injuries are most common (77.0%) in our population especially due to road traffic accidents (67.0%). Surgical management has a pivotal role in saving life where the patient is haemodynamically unstable.

Key Words: Liver Injury

Citation of articles: Bukhari AS, Hussain S, Hamza M. Evaluation of liver Injuries in DHQ Teaching Hospital Bannu. Med Forum 2017;28(12):35-38.

INTRODUCTION

Liver trauma has the potential for extensive injuries which must be carefully visualized and investigated to ensure proper management. There have been significant developments as well as changes in the management of liver trauma since last 30 years.¹ Although it is certain that most penetrating injuries require intervention, there is no absolute decision regarding the management of blunt injuries. Earlier as two decades back most blunt injuries were treated surgically to ensure homeostasis and prevent the risk of biliary leaks and sepsis. In majority of cases, the injured liver cease to hemorrhage without any intervention and a conservative approach is relatively safe in haemodynamically stable patients.

¹. Department of Surgery / Pharmacology² / Forensic Medicine³, Bannu Medical College Bannu.

Correspondence: Dr. Ajmal Shah, Associate Professor of Surgery, Bannu Medical College Bannu.

Contact No: 0300 5761970

Email: dr.ajmalbukhari11@gmail.com

Severe hepatic injuries in unstable patients require intervention and several techniques have been designed to stop hemorrhage and repair extensive parenchymal damage.

Trauma is one of the leading causes of mortality worldwide for all age groups.¹ The liver is the largest solid abdominal organ and involves majority of metabolic functions of the body.² Despite the relative protection by overlying ribs, it is susceptible to compressive forces by means of blunt trauma that can injure the soft parenchyma.^{3,4} Motor vehicle accidents are one of the most frequent causes of traumatic hepatic injury.⁴ Major liver trauma is frequently associated with coagulopathy.⁵ The developments in diagnosis, resuscitation and advent of new surgical technique have opened a new chapter in the management of liver injuries. In the past decades the use of CT SCAN has changed the diagnostic and therapeutic approach to such injuries completely,⁶ decreasing the options for surgical intervention.

MATERIALS AND METHODS

This is a retrospective clinical study that was carried out in the Department of Surgery DHQ Teaching Hospital Bannu Pakistan from 1st January 2015 to 31st December 2016. It included 60 cases of liver trauma, due to both penetrating and non-penetrating injuries, on the basis of clinical features and fulfilling inclusion criteria, admitted through accident and emergency department. This study included adult patients both male and female. Clinical data regarding age, sex, mode and type of injuries were taken and recorded. After initial resuscitation, clinical evaluation and thorough examination, those patients who were haemodynamically stable, were admitted to the ward and managed conservatively. Ultrasound or CT SCAN was done in all cases managed conservatively. Patients, who were haemodynamically unstable, managed surgically according to grading of liver injury. Resuscitation, treatment options and outcome were recorded on a Proforma, which was specifically generated for the purpose.

Inclusion criteria included age group equals or more than 13 years of age with diagnosis of liver trauma, all the patients with penetrating and non-penetrating injury to liver, all the patients with blunt and sharp injury to liver. Exclusion criteria included all patients' less than 13 years of age, patients with pre-existing liver disease i.e. cirrhosis, tumors, hepatitis etc and patients who have previously undergone hepatic surgery.

RESULTS

Table No.1: Mode of Injury

Number of cases	Penetrating Injury	Blunt Trauma
Total 60	14 (23%)	46 (76%)

Table No.2: Type of Treatment

Number of cases	Conservative treatment	Surgical Treatment
60	16 (26.6%)	44 (73%)

Table No.3: Type of Surgical Management

Number of cases	Primary suturing + perihepatic packing with topical haemostatic agents	Primary Suturing	Perihepatic Packing
44	15(34%)	9 (20%)	20 (45%)

The incidence of liver trauma due to non-penetrating injuries was 46(76.6%) while due to penetrating injuries 14 (23.3%), In all cases of blunt injuries, 40 patients(86.9%) were due to road traffic accidents, and 6 patients(13.0%) were due to assaults. 16 patients who were haemodynamically stable, were managed

Medically with strict vital monitoring, input/output charting and repeated examinations. 44 patients who were haemodynamically unstable despite aggressive resuscitation and were managed surgically. 9 patients (20%) were treated by simple suture, 15 patients (34%) were treated by suture and perihepatic packing, and 20 patients (45%) were treated by perihepatic packing. Patient who underwent perihepatic packing were re explored after 48 hours. abdomen was washed, drains were put in and abdomen was closed permanently.

Table No.4: Post Opp Complication

No. of cases	Chest Complication (pneumonia, dry cough, productive cough,)	Jaundice	Wound infection	Burst abdomen	Biliary Fistula
	30 (50%)	4 (6.6%)	9 (15%)	2 (3.3%)	3 (5%)

Table No.5: Types of injuries

Mode of injury	No of patient	Percentage
Solitary liver injury	20	45%
Associated extra hepatic injury	24	55%

DISCUSSION

One of the largest solid organ in the body is liver. Although it lies in a safe area of the body, but because of its large size, it makes it an inescapable victim of injury when the abdomen encounters like some traumatic assault. The exact incidence of liver injury is difficult to estimate due to lack of trauma registry in our country, but surely, it is high among our population. The same has been reported not only in our study, but other published local studies from different parts of the country.⁷⁻⁹ In the current study we observed predominant involvement of males. More frequent involvement of males has also been reported in some other studies in the context of trauma in general as well as liver injuries.^{7,8-10} The males are more prone to injury because they are more involved in driving, traveling and other outdoor and high-risk activities and because of that they are more frequent victims of assaults such as firearm injuries and stabs resulting from fights and brawls.

In our study, out of 60 patients, 57 patients were male and 03 were female with a male to female ratio of 19:1, this shows high male to female ratio compared to other studies because females are mostly non-Ambulant. A study conducted in Qatar by Faramawy et al, demonstrated male to female ratio of 11.6:1.¹¹ The reason for this high percentage of male preponderance is that males are more exposed and projected to trauma than females mainly due to male dominated society and outdoor activities. Similarly, in another study conducted in Saudi Arabia by Barrimahet et al, showed almost twice the number of male to female having road traffic

accidents (major cause of traumatic liver injury) in that year.

Our study found more frequent involvement of the younger population. The same has been found in other studies as well¹²⁻¹⁴. It is also reported that involvement of younger males predominantly, reflects the socio-economic implications in such injuries. In our study the predominant mechanism of liver injury was blunt, particularly RTAs. Firearm injuries as the predominant mechanism of injury is also reported in Russia and Peshawar^{14,15}. As Hemodynamic compromise ended with resuscitation and emergency exploratory laparotomy, because unacceptable delay in transportation, patients with life-threatening injuries of higher grade mostly die on the spot or on route to hospital¹⁶. In our study, in majority of patients we did perihepatic packing and suture hepatorrhaphy procedures. This is an agreement to other reported studies¹³⁻¹⁹. In fact, perihepatic packing has been great efficacy particularly in patients having liver trauma, and in conditions where blood is not available or in the cases where there is those massive transfusion requirement.

In our study, We strictly followed the policy of putting no more than six abdominal sponges around the liver, so as to avoid the complications of iatrogenic abdominal compartment syndrome. besides this, a variety of other procedures have been employed for liver injury by various researchers with reasonable good success rates. For instance, application of topical haemostatic agents, hepato-omentorrhaphy, tractotomy with finger fracture, extensive hepatorrhaphy, resectional debridement with selective vascular ligation, intra hepatic balloon, angio embolisation, venovenous bypass, and hepatic transplant etc.²⁰⁻²³ Grade of liver injury, expertise of the surgeon, preference of individual surgeon and institutional practices are the different factors which effect the haemostatic measure. Hence, standardization as well as comparison of the surgical procedures reported by various researchers is difficult to made.

In our study 47 patients (78%) presented with blunt trauma and 13 patients (21.6%) presented as penetrating while in other study 87 patients with hepatic injuries from January 1995 to December 1999. Out of which, 76% of them had sustained blunt trauma while, 24% had penetrating trauma²⁴.

In the present study 60 patients were included, out of which 20(45%) patients were with hepatic trauma alone and 24 (55%) patients were of liver trauma with associated injuries while in another study the associated complication were 62 (54.86%) patients had associated injuries⁵. in another study from LRH Peshawar the isolated liver injury were 32.5% while associated liver injuries were 67.5%.

In our study 28% patients were found to be haemodynamically stable, and managed medically with strict vital monitoring, input/output charting and repeated examination to assess the conditions. 73% patients were haemodynamically unstable despite

aggressive resuscitation and therefore were managed surgically. The choice of different surgical options for securing homeostasis in liver trauma depended on type and mode of injury, grade of liver injury and surgeon own discretion. Most of liver injuries required simple suture ligation. Simple suture ligation was done in 9 patients (20%), 20 patients were managed by perihepatic packing and 15 patients were managed by perihepatic packing and suturing. while in another study the primary suturing was in done 23% patient while perihepatic packing was done in 45% patient²⁵.

There were 40% patients in shock at the time of presentation with increase respiratory rate, pulse rate and decrease blood pressure. These patients were aggressively resuscitated with crystalloids, colloids and blood products, and shifted to emergency operation theatre for surgery.

The medical management has been time tested for haemodynamically stable patients of blunt injuries of liver. However, in penetrating injuries, especially in firearm injuries, but still exploratory laparotomy has the choice of management for many years¹⁷. This method has been objected in current studies that support selective medical treatment in haemodynamically stable cases without associated abdominal injuries. haemodynamically stable Penetrating injured patients having no signs of peritonitis, are recommended to undergo a contrast enhanced CT scan of the abdomen. If any signs shows on ultrasonography like hollow viscous perforation or evolving hemodynamic instability, exploratory laparotomy is option for treatment. The grade of the hepatic injury is not contraindicated for conservative management^(17,18).

Given the evidence base, we should evolve regulations and measures to prevent RTAs, thereby reducing the frequency of liver trauma. As majority of our patients have hemodynamic compromise and present late, morbidity and mortality rate can be decreased in our set ups due to good Advance life care trauma support. Public awareness on the issue is imperative.

in our study the chest complication is 49%, while in the study of Imran Ahmad was 44.26%,²⁶ burst abdomen in our study was 3.3% while it was reported as 1.76% in the study, in our study billiry fistula was reported in 5% patients while in study of LRH Peshawar it has been reported 5% of bile leak. wound infection was reported in our study is 3.3% while reported 10% in LRH Peshawar study. It shows that the results of our study are almost similar to the other reported findings in a periphery teaching Hospital.

CONCLUSION

In conclusion, non-penetrating liver injuries are most common (78.1%) due to road traffic accident and assault. Hemorrhage is the leading cause of death in liver trauma. In haemodynamically stable patients, non-operative management is safe and rewarding. Surgical intervention was found to be life saving in thermodynamically unstable patients.

The common surgical procedure offered is perihepatic packing and mattress suturing.

Author's Contribution:

Concept & Design of Study: Ajmal Shah Bukhari
Drafting: Shabir Hussain and Mir Hamza

Data Analysis: Shabir Hussain and Mir Hamza

Revisiting Critically: Ajmal Shah Bukhari, Shabir Hussain and Mir Hamza

Final Approval of version: Ajmal Shah Bukhari

Conflict of Interest: The study has no conflict of interest to declare by any author.

REFERENCES

- Mahran DG, Farouk OA, Qayed M, Berraud AF. Hospitalized injuries and deaths in a trauma unit in upper Egypt. *Int J Critical Illness Injury Sci* 2013; 3(4):235-40.
- Munding J, Tannapfel A. Anatomie der Leber: Was muss der Radiologe wissen? *Der Radiologe* 2011;51(8):655-60.
- Chien L-C, Lo S-S, Yeh S-Y. Incidence of liver trauma and relative risk factors for mortality: A population-based study. *J Chinese Med Assoc* 2013;76(10):576-82.
- Saaq M, Niaz-ud-Din MZ, Shah SA. Presentation and outcome of surgically managed liver trauma: experience at a tertiary care teaching hospital. *JPMA* 2013;63(4):436-9.
- Armand R, Hess JR. Treating coagulopathy in trauma patients. *Transfusion Medicine Reviews* 2003;17(3):223-31.
- Hassan R, Aziz AA. Computed Tomography (CT) Imaging of Injuries from Blunt Abdominal Trauma: A Pictorial Essay. *Malaysian J Med Sci* 2010;17(2):29-39.
- Muizuddin M, Rehman A, Alam SN, Manzar S. Intra-abdominal visceral injuries in blunt abdominal trauma.
- Khan JS, Iqbal N, Gardezi JR. Pattern of visceral injuries following blunt abdominal trauma in motor vehicular accidents. *J Coll Physicians Surg Pak* 2006; 16: 645-7.
- Ali U, Noor A, Shah MM, Alam W. Trauma management in a tertiary care hospital in Peshawar, Pakistan. *J Ayub Med Coll Abbottabad* 2008;20: 112-6.
- Saaq M, Shah SA. Thoracic trauma: Presentation and management outcome. *J Coll Physicians Surg Pak* 2008;18:230-3.
- El-Faramawy A, El-Menyar A, Zarour A, Maull K, Riebe J, Kumar K, et al. Presentation and outcome of traumatic spinal fractures. *J Emerg Trauma Shock* 2012;5(4):316-20.
- Butt HA, Zahur S, Haq AU. Hepatic trauma: effects of delay in presentation. *Ann King Edward Med Uni* 2006;12: 293-5.
- Khan JS, Iqbal N, Gardezi JR. Pattern of visceral injuries following blunt abdominal trauma in motor vehicular accidents. *J Coll Physicians Surg Pak* 2006;16:645-7.
- Ali U, Noor A, Shah MM, Alam W. Trauma management in a tertiary care hospital in Peshawar, Pakistan. *J Ayub Med Coll Abbottabad* 2008; 20: 112-6.
- Urman MG, Subbotin AV. Surgical strategy for the liver injury and developed abdominal complications. *Vestn Khir Im II Grek* 2009;168: 72-5.
- Saaq M, Shah SA. Thoracic trauma: Presentation and management outcome. *J Coll Physicians Surg Pak* 2008;18:230-3.
- Saaq M, Shah SA. Thoracic trauma: Presentation and management outcome. *J Coll Physicians Surg Pak* 2008;18:230-3.
- Vatanaprasan T. Operative treatment of hepatic trauma in Vachira Phuket hospital. *J Med Assoc Thai* 2005;88: 318-28.
- Gao JM, Du DY, Zhao XJ, Liu GL, Yang J, Zhao SH, et al. Liver trauma: experience in 348 cases. *World J Surg* 2003; 27: 703-8.
- Ahmed N, Vernick JJ. Management of liver trauma in adults. *J Emerg Trauma Shock* 2011; 4: 114-9.
- Badger SA, Barclay R, Campbell P, Mole DJ, Diamond T. Management of liver trauma. *World J Surg* 2009; 33: 2522-37.
- Mohr AM, Lavery RF, Barone A, Bahrapour P, Magnotti LJ, Osband AJ, et al. Angiographic embolization for liver injuries: low mortality, high morbidity. *J Trauma* 2003; 55: 1077-81.
- Delis SG, Bakoyiannis A, Selvaggi G, Weppler D, Levi D, Tzakis AG. Liver transplantation for severe hepatic trauma: Experience from a single center. *World J Gastroenterol* 2009; 15: 1641-4.
- Ratta plee Pak-art, ChadinTharavej Boonchoo Sirichandkul A multidisciplinary approach in the management of hepatic injuries. *Injury* 2002;33(4): 309-315.
- The General Surgery and Trauma Service of the Hospital da Restauração – SUS, Recife, PE, Brazil 2013;26(2):129-132
- Ahmad I. Hepatic trauma and associated injuries - Experience in a tertiary care hospital. *Pakistan Oral Dental J* 2014;34(2): 243-244.