

# Peripheral Neuropathy: Incidence and Clinical Presentation in the Cases of Diabetic Mellitus

1. Muhammad Iqal 2. Abdul Munam 3. Adnan Ahmed

1. Assoc. Prof. of Medicine, LUHMS, Jamshoro 2. Assoc. Prof. of Neurology, Isra University Hospital, Hyderabad  
3. Asstt. Prof. of Radiology, LUHMS Jamshoro

## ABSTRACT

**Objective:** To assess the incidence and clinical presentation of peripheral neuropathy in the cases of diabetic mellitus

**Study Design:** Cross sectional study

**Place and Duration of Study:** This study was conducted at Medicine Department of LUMHS with the duration of time February 2011 to January 2012.

**Materials and Methods:** Total 200 cases were chosen for this study after diagnosis of diabetic mellitus. All equitable laboratory investigations were carried out, along with HbA1C. All the cases were excluded from the study those were less than 5 HbA1c levels, history of any trauma leading to neurological deficit that can compromise assessment of sensory system along with history of renal failure. All the data regarding demographic characteristics, incidence and clinical pattern of the cases with peripheral neuropathy was entered in the pre-designed Proforma.

**Results:** Mean age of the cases was  $56.3 \pm 10.21$  years. Mostly cases were noted from urban area 60.0%. 46.0% cases were smoker and mostly smoker were male, while 26(13.0%) patients (totally male) were with history of alcohol consumption. Mostly patients were poor socioeconomically. Incidence of the neuropathy was noted 60% further according to the classification, 16% cases were found with mild neuropathy, 21% were with severe while majority of the cases 32% were found with moderate neuropathy. According to the clinical presentation mostly cases were found with burning, 85.0%, Pain 66.6%, numbness 75.0%, while Sensitive touch, Itching, Foot ulcer, Muscle cramp and Tingling were found as 60(50.0%), 26(30.0%), 18(15.0%), 45(37.5%) and 51(42.5%) respectively.

**Conclusion:** In this study it is concluded that incidence of diabetic peripheral neuropathy is very high, and it increases with the increasing time of diabetes, uncontrolled diabetes and ignorance diabetes. Majority of the cases had found with sensory loss having foot ulcers, and they were unaware regarding this condition.

**Key Words:** Peripheral Neuropathy, Clinical Presentation, Diabetic Mellitus

**Citation of article:** Iqal M, Munam A, Ahmed A. Peripheral Neuropathy: Incidence and Clinical Presentation in the Cases of Diabetic Mellitus. Med Forum 2015;26(8): 47-50.

## INTRODUCTION

Diabetes mellitus (DM) has come to pandemic extents around the world. Generally, diabetes was viewed as a big problem kept to created nations and rich individuals. Nonetheless, newly gauges propose that incidence of diabetes is increasing all around, especially in creating nations.<sup>1</sup> DM has turned into an critical wellbeing concern in South Asian along with an expected increment in the incidence more than 15.1% somewhere around 2000 and 2030.<sup>1</sup> Neuropathy is viewed as the most widely recognized micro vascular risk for in type I and II DM.<sup>2,3</sup> Neuropathic issue in diabetes can damage function of CNS, PNS and/or ANS.<sup>4</sup> Distal peripheral neuropathy (DPN), called diabetic polyneuropathy influences PNS and is by a

wide margin the most well-known sort of neuropathy found in DM. In results functioning loss of peripheral nerves causes loss of defensive sensations and damage ability of the cases to see early or even obvious feet ulcerations. DPN is viewed as a principle reason of the amputation, and subsequently a huge reason for morbidity in diabetic mellitus.<sup>5</sup>

In spite of the fact that a typical and essential diabetic complication, neuropathy has not been concentrated on as frequently or as widely complication of macro vascular, retinopathy or nephropathy. What's more, the reported predominance estimated broadly between nations, to some degree because of the distinction in testing techniques and absence of accord on criteria of diagnosis.<sup>6</sup> Rising age, longer length of time of diabetes and diabetic poor control are very much perceived danger elements for DPN, while retinopathy, hypertension, weight, hyperlipidaemia, smoking, and microalbuminuria have additionally been ensnared as potential danger markers.<sup>7</sup> Most commonness and danger variable studies are from western created

**Correspondence:** Dr. Muhammad Iqal,  
Assoc. Prof. of Medicine, LUHMS, Jamshoro  
**Cell No.:** 0333-2700190  
**E-mail:** dr.saeedarain786@gmail.com

nations, while a linked data shortage from creating nations, especially from the region of the South-Asian.<sup>8</sup> In any case, an expected 80% of the worldwide populace with diabetes lives in creating nations. Population of the South-Asia specifically is well known with an expanded inclination for the illness. What's more, a few studies have shown that the danger of diabetes related removals and the incidence of ulcer of the diabetic foot UK is fundamentally lower in cases of the diabetes of Asian inception when contrasted with that of diabetic cases of European root.<sup>9,10</sup> This diminished risk in Asians was observed to be identified with the lower occurrence of peripheral artery and DPN. Ethnic contrasts and differential ecological presentation to hazard components in the various nations are extra proposed systems. This study was conducted to see the incidence and clinical presentation of peripheral neuropathy in the patients with diabetic mellitus.

**MATERIALS AND METHODS**


This cross sectional study was performed at medicine department of LUMHS with the duration of time February 2011 to January 2012. Total 200 cases were chosen in this study after diagnosis of diabetic mellitus, all the cases were selected above the age 30 years. All the cases with diabetic mellitus those were agree to participation were preferred. Full medical history of the cases with complete physical examinations was carried out. All equitable lab investigations were carried out along with HbA1c. All the cases were excluded from the study those which were less than 5 HbA1c level, any trauma leading to neurological deficit that can compromise assessment of sensory system along with history of renal failure. Neuropathy degrees were done on the basis of TCSS score utilized in the previous study as {no neuropathy: ≤5, mild: 6-8, moderate: 9-11 and severe: ≥12}.<sup>11</sup> All the data regarding demographic characteristics, incidence of and clinical presentation of the cases with peripheral neuropathy was entered in the predesigned Performa.

**RESULTS**

Total 200 known diabetic cases were selected in the study; patients mean age was 56.3 + 10.21 years. High difference was not found in male and female, as male 104(52.0%) and female 96(48.0%). Mostly cases were noted from urban area 120(60.0%), while 80(40.0%) cases were comes from rural areas. 92(46.0%) cases were smoker and mostly male, while 26(13.0%) patients (totally male) were with history of alcohol consumption. Mostly patients were poor socioeconomically. **Table 1.** 16% cases were found with mild neuropathy, 21% were with severe while majority of the cases 32% were found with moderate neuropathy. **Figure 1**

**Table No. 1: Demographic characteristics of patients (n = 200)**

Basic status	No. of patients/percentage
Age (Mean+SD)	56.3 + 10.21
<b>Gender</b>	
Male	104(52.0%)
Female	96(48.0%)
<b>Residence</b>	
Rural	80(40.0%)
Urban	120(60.0%)
<b>Smoking status</b>	
Yes	92(46.0%)
No	108(54.0%)
<b>Alcohol consumption</b>	
Yes	26(13.0%)
No	124(58.0%)
<b>Obesity</b>	
Yes	37(18.5%)
No	163(81.5%)



**Figure No.1: Incidence of Peripheral Neuropathy. n=200**

**Table No. 2; Presentation of the cases with PN n=120**

Presentation	Frequency/percentage
Burning	102(85.0%)
Pain	80(66.6%)
Sensitive touch	60(50.0%)
Numbness	90(75.0%)
Itching	36(30.0%)
Foot ulcer	18(15.0%)
Muscle cramp	45(37.5%)
Tingling	51(42.5%)

According to the clinical presentation mostly cases were found with burning, 102(85.0%), Pain was found in 80(66.6%), numbness was found in 90(75.0%), while Sensitive touch, Itching, Foot ulcer, Muscle cramp and Tingling were found as 60(50.0%), 36(30.0%), 18(15.0%), 45(37.5%) and 51(42.5%) respectively. **Table 2.**

## DISCUSSION

Diabetes one of the common cause of different complications in the patients as renal disease, blindness and lower leg amputation even though the occurrence of different treatment approaches these all supply to the excess morbidity as well as mortality in the cases with diabetes.<sup>12</sup>

In this series mostly cases were found with old age above 40 years, only few cases were noted with age of less than 40, as well as mean age of the cases was found mean+SD 56.3 + 10.21. Similarly Iftikhar M. et al<sup>13</sup> reported mean age of the cases as mean+SD 49.52±7.933. Salzedas Muniz EC et al<sup>14</sup> reported mean age as 60.9 years. Al-Sarihin K et al<sup>15</sup> mentioned mean age of the cases as 56.19±14.31 years.

High difference was not found in male and female, as male 104(52.0%) and female 96(48.0%).

Mostly cases were noted from urban area 120(60.0%). 92(46.0%) cases were smoker and mostly smokers were male. A.B. Zafar et al<sup>16</sup> stated 49 male and 50 female in his study. Iftikhar M. et al<sup>13</sup> and Al-Sarihin K et al<sup>15</sup> also reported similar findings of gender, residence and smoking status.

PDN is probably main complication that's affects are concern to sensory, autonomic, and motor neurons of the peripheral nervous system. WHO reported that about fifty percent in diabetic patients develop neuropathy.<sup>17</sup> With increase time of diabetic, DN risks also increase. It is reported that 60 to 70% neuropathy developed in cases with diabetes.<sup>18</sup> Iftikhar M. et al<sup>13</sup> stated incidence of DN in diabetic patients was 127 (74.8%). Feray Soyupek *et al*<sup>19</sup> mentioned that in the study frequency of peripheral neuropathy was 80.4%. While in this study 60% patients were found with peripheral neuropathy, out of them 16% cases were found with mild neuropathy, 21% with severe while great part of the cases 32% were found with moderate neuropathy, similarly Davies M. et al<sup>20</sup> found neuropathy as mild 20%, moderate 47%, or sever 33%. Clinical pattern DPN is associated that incorporates troublesome pain toward one side and there may be unfeeling foot on the flip side that likewise at danger of ulceration. In the advanced PDN the raised vibration and warm discernment edges that prompts degeneration and the sensory loss of total categories in peripheral nerve of the fibers. According to the clinical presentation in the present series out of 120 patients mostly cases were found with burning, 102(85.0%), Pain was found in 80(66.6%), numbness was found in 90(75.0%), Sensitive touch 60(50.0%), Itching 36(30.0%), Muscle cramp 45(37.5%) and Tingling were found in 51(42.5%) patients. Iftikhar M. et al<sup>13</sup> stated that Out of 250 patients 116 (46.4 %) cases noted with burning type pain and 114 (45.6%) were found to have numbness while 61 (24.4%) had tingling. Out of 250 patients 17 (6.8%) were found to have sharp

pain and 55 (22%) had dullness. Out of 250 patients 44 (17.6%) had coldness. Katulanda P et al<sup>21</sup> reported patients with established diabetes the most common symptom was the presence of numbness of the feet 37.1%, followed by burning, aching or tenderness of the feet 32.3%, prickling sensation 29.7% and unsteadiness 25.5%.

Initial screening of PDN is basic in light of the fact that patients of all phases of neuropathy are at a more serious danger of creating uncaring foot ulceration. The group based study directed at Karachi, Pakistan stated that DPN developed in the around 40% cases, and out of them 4% foot ulcers developed.<sup>22</sup> likely in another study consider that development of diabetic neuropathy is 36%, and foot ulcer 10.4% in diabetes cases.<sup>23</sup> As well s in the present study PDN was found 60% and 15.0% patients were found with foot ulceration..

## CONCLUSION

In this study it is concluded that diabetic peripheral neuropathy is increased with the increasing time of diabetes, causing uncontrolled and ignored diabetes and male/female are involve. Majority of the cases had found with sensory loss having foot ulcers, and they were unaware regarding this condition. Government should facilitate the diabetic patients regarding awareness and complete treatment, because in this study mostly patients were poor socioeconomically. More research is needed to better identification of factors associate to DPN.

**Conflict of Interest:** The study has no conflict of interest to declare by any author.

## REFERENCES

1. Wild S, Roglic G, Green A, Sicree R, King H: Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care* 2004; 27(5):1047–1053.
2. Abbott CA, Carrington AL, Ashe H, Bath S, Every LC, Griffiths J, Hann AW, Hussein A, Jackson N, Johnson KE, et al: The North-West Diabetes Foot Care Study: incidence of, and risk factors for, new diabetic foot ulceration in a community-based patient cohort. *Diabet Med* 2002;19(5):377–384.
3. Daousi C, MacFarlane IA, Woodward A, Nurmikko TJ, Bundred PE, Benbow SJ: Chronic painful peripheral neuropathy in an urban community: a controlled comparison of people with and without diabetes. *Diabet Med* 2004;21(9):976–982.
4. American Diabetes Association, American Academy of Neurology: Consensus statement: report and recommendations of the san Antonio conference on diabetic neuropathy. *american diabetes association american academy of neurology. Diabetes Care* 1988;11(7):592–597.

5. Jeffcoate WJ, Harding KG: Diabetic foot ulcers. *Lancet* 2003; 36:1545–1551.
6. Schmader KE: Epidemiology and impact on quality of life of postherpetic neuralgia and painful diabetic neuropathy. *Clin J Pain* 2002;18(6):350–354.
7. Boulton AJ, Cavanagh PR, Rayman G: The foot in diabetes. 4 edition. Hoboken: John Wiley & Sons Ltd; 2006.
8. Morkrid K, Ali L, Hussain A: Risk factors and prevalence of diabetic peripheral neuropathy: a study of type 2 diabetic outpatients in Bangladesh. *Int J Diabetes Dev Ctries* 2010;30(1):11–17.
9. Abbott CA, Garrow AP, Carrington AL, Morris J, Van Ross ER, Boulton AJ: Foot ulcer risk is lower in South-Asian and african-Caribbean compared with European diabetic patients in the U.K.: the North-West diabetes foot care study. *Diabetes Care* 2005;28(8):1869–1875.
10. Chaturvedi N, Abbott CA, Whalley A, Widdows P, Leggetter SY, Boulton AJ: Risk of diabetes-related amputation in South Asians vs. Europeans in the UK. *Diabet Med* 2002;19(2):99–104.
11. Perkins BA, Olaleye D, Zinman B, Bril V: Simple screening tests for peripheral neuropathy in the diabetes clinic. *Diabetes Care* 2001;24 (2):250–256.
12. Girach A, Vignati L. Diabetic microvascular complications-can the presence of one predict the development of another? *J Diabetes Complications* 2006;20:228–37.
13. Iftikhar m, Hussain a, Rizvi a. Frequency of peripheral neuropathy in patients with diabetes mellitus. *J ayub med coll abottabad* 2014; 26(4);584-86
14. Muniz EC, Rocha RM, Reis ML, Santos VL, Grossi SA. Neuropathic and ischemic changes of the foot in Brazilian patients with diabetes. *Ostomy Wound Manag* 2003; 49:60–70.
15. Al-Sarihin K, Althwabia I, Khaled M, Haddad f. Prevalence of peripheral neuropathy among patients with diabetes mellitus at King Hussein Hospital, Amman, Jordan. *Rawal Medical Journal* 2013;38;2;92-96.
16. A.B. Zafar , J. Ahmed , A. Basit , S.F.D. Alvi , A. Fawwad, M. Riaz. Factors screening the peripheral neuropathy in asymptomatic patients with type 2 diabetes by Neurothesiometer . *Journal of Diabetology*, October 2014; 3:2;1-6
17. World Health Organisation. Diabetes.[http://www.who.int/hpr/NPH/docs/gs\\_diabetes.pdf](http://www.who.int/hpr/NPH/docs/gs_diabetes.pdf). (Accessed on 01 February 2010)
18. National Diabetes Information Clearinghouse. “Diabetic Neuropathies: The Nerve Damage of Diabetes.” <http://diabetes.niddk.nih.gov/>, 2009. Accessed on 20 February, 2014.
19. Soyupek F, Ceceli E, Suslu FE, Yorgancioğlu R. Neurologic and radiologic abnormalities of the foot in diabetic patients. *J Pak Musculoskelet Rehabil* 2007;20:55–60.
20. Katulanda P, Ranasinghe P, Jayawardena R, Constantinou G, Sheriff MH, Matthews DR. The prevalence, patterns and predictors of diabetic peripheral neuropathy in a developing country. *Diabetol Metab Syndr*. 2012 May 29;4(1):21.
21. Davies M, Brophy S, Williams R, Taylor A. The prevalence, severity, and impact of painful diabetic peripheral neuropathy in type 2 diabetes. *Diabetes Care*. 2006 Jul;29(7):1518-22.
22. Jawad F, Maqsood A, Jamal S, Azfar M, Ahmed U, Shera SA. Prevalence of chronic complications and association in Type-2 diabetes. *J Pak Med Assoc*. 2004; 54: 54-59.
23. Basit A, Hydrie MZ, Hakeem R, Ahmedani MY, Masood Q. Frequency of chronic complications of type 2 diabetes. *J Coll Physicians Surg Pak*. 2004; 14:79-83.