

Outcome of Surgical and Medical Management of Anal Fissure

1. Majeed Ullah Buzdar 2. Irfan Qasirani

1. Asstt. Prof, of Surgery, 2. PGR Department of Surgery Ghazi Khan Medical College, D.G.Khan

ABSTRACT

Objective: To study the outcome of Surgical and Medical treatment of Anal Fissure.

Study Design: Prospective study

Place and Duration of Study: This was study carried out in Surgical Department of Ghazi Khan Medical College, Dera Ghazi Khan, from January 2014 to December 2014.

Materials and Methods: A total of 75 patients form Surgical OPD were included in the study. Patients having anal fissure were diagnosed clinically and were selected according to inclusion criteria.

Results: Out of 75 patients, 49 (65%) were male and 26 (35%) were female patients. The patients were from the ages of 23 to 54. Painful defecation was present in all the patients (100%), constipation in 67 (90%) patients, whereas bleeding per rectum in 52(70%). Sentinel pile was seen in 50 (67%) patients and associated superficial fistula only in 1 patient.

Conclusion: it is concluded that lateral internal sphincterotomy is the most effective way of treatment of chronic anal fissure, whereas chemical sphincterotomy with topical glyceryl trinitrate is relatively less effective.

Key Words: Dentate line, lateral sphincterotomy, defecation

Citation of article: Buzdar MU, Qasirani I. Outcome of Surgical and Medical Management of Anal Fissure. Med Forum 2015;26(9):42-44.

INTRODUCTION

Anal Fissure is a linear tear or ulceration in the lining of the squamous epithelium in the anal canal distal to dentate line (mucocutaneous junction) due to local trauma. The fissure causes excruciating pain during defecation that persists for two to three hours. Hypertonicity and hypertrophy of the internal sphincter is so severe that it causes spasm, pain and ischemia leading to non-healing of the fissure¹.

The fibers of the internal anal sphincter are visible in the base of chronic anal fissure and often an enlarged anal skin tag is present in the anal canal proximal to the fissure known as sentinel pile.

The lateral internal sphincterotomy is the first line surgical option for all the fissures associated with hypertonicity and hypertrophy of the internal anal sphincter^{3,4,5}. It can be performed using open or closed methods, depending upon the surgeon's choice.

Another procedure by advancement flap, is usually reserved for recurrent fissures or fissures with low pressure (Tone). The procedures, like Lord's anal dilatation and mid-line posterior sphincterotomy, are obsolete because of high rates of recurrence, incontinence and delayed wound healing^{6,7}.

The conventional treatment of anal fissure is surgical lateral internal sphincterotomy. The alternative option of Chemical sphincterotomy using medication such as

topical glyceryl trinitrate induces rapid healing of anal fissure. It is a new, easily handled and affective alternative to surgical lateral sphincterotomy⁸. It offers a significant healing rate for acute anal fissure and prevents it's evolution to chronicity⁹.

MATERIALS AND METHODS

This was a prospective study carried out in Surgical department of Ghazi Khan Medical College, Dera Ghazi Khan, from January 2014 to December 2014. A total of 75 patients form Surgical OPD were included in the study. Patients having anal fissure were diagnosed clinically and were selected according to inclusion criteria. All patients were divided in two groups, group A patients were treated surgically undergoing lateral internal sphincterotomy and group B patients were treated medically with topical 0.2% glyceryl trinitrate and adjuncts.

RESULTS

Among 75 patients, 49 (65%) were male and 26 (35%) were female.

Table No.1: Gender Distribution (N=75)

Gender	No. Of patients	Percentage
Male	49	65%
Female	26	35%

All the patients were between the ages of 23 to 54 year.

Regarding symptoms, painful defecation was present in all the patients (100%), constipation was a feature in 67 (90%) patients while bleeding per rectum was documented in 52 (70%) patients. Sentinel pile was

Correspondence: Dr. Majeed Ullah Buzdar
Asstt. Prof, of Surgery, Ghazi Khan Medical College,
D.G.Khan
Cell No.: 0334-7473540
E-mail: ullahmajeed@yahoo.com

found in 50 (67%) of the patients whereas associated superficial fistula was noted only in 1 patient.

Table No.2: Age Distribution (n=75)

Age Years	No of Patient	Percentage
20—30	32	40%
30—40	26	36%
40—50	13	18%
>50	4	6%

Table No.3: Presentation (symptoms) (n=75)

Symptoms	No. of Patients	Percentage
Painful defecation	75	100%
Constipation	62	90%
Bleeding per rectum	50	70%
Associated fistula	1	1.3%

Anal Fissure was present at the posterior mid-line in 61 (82%) patients, in 9(12%) patients fissure was lateral in position while in 5(6%) patients, it was present at anterior mid-line.

Table No.4: Position of Anal Fissure (n=75)

Position	No. of Patients	Percentage
Posterior Midline	60	82%
Lateral	9	12%
Anterior Midline	5	6%

Out of 75 patients, 48(64%) opted for surgical treatment and underwent lateral internal sphincterotomy, were placed in group A whereas 27(36%) patients opted for medical treatment with topical glyceryl trinitrate and adjuncts, were placed in group B.

Table No.5: Treatment option (n=75)

Treatment Option	No. of Patients	Percentage
Surgical	48	64%
Medical	27	36%

DISCUSSION

All the patients were given the surgical and medical options of the treatment and were placed in group A or B according to their selected choice after informed consent. In our study, 48(64%) of the patients who opted for Surgical treatment underwent lateral internal sphincterotomy and all of them had uneventful recovery in a short period of time but one of them had some soiling. Our results are comparable to Lysy, who had 100% healing and 0% recurrence with lateral internal sphincterotomy¹⁰.

Though, surgery for anal fissure is associated with few complications like permanent incontinence of faeces, transient in-continnence of flatus and soiling but such complications can be prevented by the use of judicious

surgical techniques and by familiarity with ano rectal anatomy. Gosse link found a flatus in-continnence rate in 30% of the patients after lateral internal sphincterotomy¹¹, in-another retrospective study where the patients underwent closed or open lateral internal sphincterotomy, 21% cases had a flatus or liquid incontinence¹².

In our study, 27(36%) patients who opted for medical treatment underwent chemical sphincterotomy with topical 0.2% glyceryl trinitrate, 20(74%) of the were cured over a period of 06 weeks while 7(26%) of the patients had a prolonged duration of healing and 2 of them opted to undergo the Surgical treatment. Our result are comparable to some other studies who have shown healing rate up to 70% with topical glycerin trinitrate^{13,14,15}.

Comparing the efficacy of surgical and medical options on the basis of effectiveness regarding the control of symptoms and side effects, both are comparable, though topical modality has a relatively higher recurrence and persistence rate but with insignificant side-effect like headache. Whereas, the surgical option, lateral internal sphincterotomy is associated with serious side-effects like permanent in-continnence of faeces or flatus and soiling.

CONCLUSION

It is concluded that in patients with anal fissure, lateral internal sphincterotomy is the state of art gold standard procedure but associated with some serious side-effects, while chemical sphincterotomy with topical glyceryl trinitrate is a suitable and reliable alternative to lateral internal sphincterotomy which can be offered as a first line treatment to the patients presenting with anal fissure.

Conflict of Interest: The study has no conflict of interest to declare by any author.

REFERENCES

1. Haq Z, Rehman M, Choudhry RA. Chemical sphincterotomy- First line of treatment for chronic anal fissure. *Mymensing Med J* 2005; 14:88-90.
2. Scholefield JH, Bock JU, Maria B. A dose finding study with 0.1% and 0.4% glyceryl nitrate ointment in patients with chronic anal fissure. *GUT* 2003; 52:264-9.
3. Herzig Do, Lu KC. Anal fissure. *Surg Clin N Am* 2010; 90(1):33-44.
4. Orsay C, Rakinic J, Perry WB, Hyman N, Buie D, Cataldo P, et al. Practice Parameters for the management of anal fissures. *Dis Colon Rectum* 2004; 47(12):2003-7.
5. Richard CS, regoire R, Plewes EA, Silverman R, Burul C, Buie D et al. Internal sphincterotomy is superior to topical nitroglycerin in the treatment of

- chronic anal fissure. Dis colon Rectum 2000; 43(8): 1048.
6. Aziz R, Din F, Shoaib M, Non-surgical treatment of chronic anal fissure. Ann KE Med Coll 2005; 11:396-7.
 7. Abd Elhady HM, Othman IH, Hablus MA, et al long-term prospective randomized clinical and manometric comparison between surgical and chemical sphincterotomy for treatment of chronic anal fissure. S Afr J Surg 209; 47(4): 112-4.
 8. Simpson J, Lund JN, Thompson RJ. The use of GTN in the treatment of chronic anal fissure. Med Sci Monit 2013; 9:123-6.
 9. Lysy J, Israeli E, Levy S. Chemical sphincterotomy first line treatment for chronic anal fissure. Dis Colon Rectum 2009; 49:858-64.
 10. Mousavi SR, Sharifi M, Medikhah Z. A Comparison between the results of fissurectomy and lateral internal sphincterotomy in the surgical management of chronic and fissure. J Gastrointest Surg 2009; 13(7):1279-82.
 11. Nelson R. A Systematic review of medical therapy for anal fissure. Dis Colon Rectum 2004; 47(4): 4422-31.
 12. Gamer Jp, McFall M, Edwards DP. J R. The medical and surgical management of chronic anal fissure. Army Med Corps 20; 148(3): 230-5 2012.
 13. Kocher HM, Steward M, Leather AJ, Cullent PT Randomized Clinical trial assessing the side effects of glyceryl trinitrate and diltiazem hydrochloride in the treatment of chronic anal fissure. Br J Surg 2002; 89:413-7.
 14. Schiano di visconte M, Munegato G. Glyceryl trinitrate ointment (0.25%) and anal cryothermal dilators in the treatment of chronic anal fissures. J Gastrointest Surg 2009; 13(7): 1283-91.
 15. Novell F, Novell CF, Novell J. Topical glycerol trinitrate in the treatment of anal fissure. Rev Esp Enterm Dig 2004; 96:255-8

Electronic Copy