

Ocular Injuries - An Experience at Anwar Paracha Eye Hospital Sukkur

1. Muhammad Afzal Pechuho 2. Kanwal Kumar 3. Rasheed Ahmad

1. Assoc. Prof. & HOD Ophthalmology, 2. Assoc. Prof. of Forensic Medicine, 3. Asstt. Prof. of Forensic Medicine, Ghulam Mohammad Mahar Medical College, Sukkur.

ABSTRACT5.

Objective: To provide the spectrum of ocular trauma, their potential hazards and propose preventive measures.

Study Design: Retrospective Descriptive

Place and Duration of Study: This study was conducted in the departments of Anwar Paracha Eye Hospital (APEH) & Forensic Medicine Ghulam Mohammad Mahar Medical College Sukkur (GMMMC) & Chandka Medical College Larkana (CMCL), constituent colleges of Shaheed Mohtarma Benazir Bhutto Medical University Larkana (SMBBMUL) from January to December 2013.

Materials and Methods: The present study was conducted from the records of patients of ocular injuries who attended Anwar Paracha Eye Hospital GMMMC Sukkur with association of departments of Forensic Medicine of GMMMC Sukkur & CMC Larkana, the constituent colleges of SMBBMUL.

Results: Of all patients visited APEH, 1.016% cases were of ocular injuries with Male to Female ratio of 8.34:1. The most vulnerable age group was 11-30 years (67.14%). On presentation in most of cases 130 (46.43%) had visual acuity 6/9 & on follow up 6/6 in 200 (71.43%) cases. The mainly sharp objects in 200 (71.43%) cases were to cause injury and mostly manner of injury was an accident in 212(75.72%) cases.

Conclusion: Majority of the cases were male of young age injured by sharp object due to accident with good prognosis of visual acuity.

Key word: Ocular, Injury, Frequency, Accidents

INTRODUCTION

In our body eyes are in prominent location, though protected by the blinking lids, orbital bones and cushioned retro-bulbar fat but more prone to trauma by variety of substances specially at workplace in many life¹. Every year more than half a million ocular injuries causing blindness worldwide that restrict activities for more than a day and approximately 19 million have unilateral permanent reduction in sight at least & 1.6 million people are blinded because of ocular injuries². The variety of articles that may be put to use as weapon like knife, razor, stick, stone, needle dagger, icpick, fire-arm³. The weapon is described as any article made or adapted for use to cause injury to the person⁴. The weapons causing eye injuries are categorized into blunt, sharp/penetrating and foreign bodies. A trauma produced by blunt weapon to the eye may result in a spectrum of injuries ranging from a simple 'Black eye' to traumatic mydriasis, iridodialysis, hyphaemia, concessional cataract, lens subluxation or complete dislocation, retinal dialysis, retinal oedema & haemorrhage, vitreous haemorrhage, a retinal blood vessel injury by compressing antero-posteriorly & stretching in the equatorial plane correspondingly². Sharp/Penetrating eye injuries depend on where & how for the object enters into the eye, Isolated to cornea or corneo-sclera, Anterior & posterior segments of eye ball. Foreign bodies as they enter & pass through the eye if not removed rapidly (1) may cause toxicity to

tissues as they degrade or oxidize or (2) damage to the intraocular contents.

The rationale is that the eye injuries are generally considered in the context of clinical eye care delivery system and not awareing the public of their potential hazards and preventive strategies commonly, so the present study is an attempt to address this deficit in this zone by providing the profile of ocular injury cases.

MATERIALS AND METHODS

The present study was conducted from the records of patients of ocular injuries who attended Anwar Paracha Eye Hospital GMMMC Sukkur with association of departments of Forensic Medicine of GMMMC Sukkur & CMC Larkana, the constituent colleges of SMBBMUL from January to December 2013. The variables considered gender, age, type of weapon, visual acuity assessed by Snellien chart & manner of injuries were entered in Statistical Package of Social sciences (SPSS) version-17. Findings were expressed in numbers & percentage. This study was approved by the Ethical Review Committee of Shaheed Mohtarma Benazir Bhutto Medical University Larkana.

RESULTS

A total 27542 patients visited to eye Hospital, of which 280 (1.016%) were enrolled of ocular injuries as shown in graph No: 1. Out of 280, males were 250 (89.29%) and females 30 (10.71%) with M:F ratio 8.34:1 as shown in graph No: 2. Majority of the patients 106

(37.86%) were between 21-30 years of age followed by 82 (29.28%) patients in 11-20 years age group and the least number of cases 07 (2.50%) in age group of 61 & above years as shown in table No: 1.

Table No. 1: Age distribution (n=280)

Age group	No	%age
Less than 10 years	25	8.93
11- 20 years	82	29.28
21-30 years	106	37.86
31-40 years	26	09.28
41-50 years	24	08.57
51-60 years	10	3.57
61 & above	07	2.5
Total	280	100%

Table No. 2: Co-relation of kind of weapon to type of injury. (n=280)

Type of injury	Kind of weapon	No	%	Total (%)
sharp	Scissor	77	27.5	200 (71.43%)
	Knife	42	15.00	
	Screw driver	30	10.71	
	Used syringe needle	16	05.71	
	Wind screen glass	12	04.29	
	Iron rod	10	03.57	
	Tip of ball point pen	05	01.79	
	Air gun	04	01.43	
	Kite stick	04	01.43	
Blunt	Stone	12	04.29	50 (17.43%)
	Hand bell	10	03.57	
	Tennis ball	08	02.86	
	Gili danda	06	02.14	
	Edge of brick	04	01.43	
	Tree twig	02	00.71	
	Door handle	02	00.71	
	Unknown	06	02.14	
Foreign body	Cracker	09	03.21	30 (10.71%)
	Explosive material	07	02.50	
	Air gun pellets	04	01.43	
	Metallic foreign body	10	03.57	
Total		280	100%	280 (100%)

Sharp objects were the most common 200 (71.43%) followed by blunt 50 (17.86%) and the least involvement by Foreign bodies 30 (10.71%) to cause the ocular injuries as shown in table No: 2. The visual acuity on presentation was as; in patients 90 (32.14%) 6/6, 130 (46.43%) 6/9, 12 (4.29%) 6/18, 18 (6.43%) 6/36, 23 (8.21%) 6/24, 05(1.79%)<6/60 and 02 (0.71%) PI +ve on Snellen Chart while on final presentation had visual acuity as; 200 (71.43%) 6/6, 46 (16.43%) 6/9, 04 (1.43%) 6/18, 07 (2.5%) 6/36, 17 (6.07%) 6/24, 04 (1.43%) <6/60 & 02 (0.71%) +ve as shown in table NO. 3. The manner of injuries was accidental in


212 (75.72%) and 68 (24.28%) homicidal as shown in table No. 4.

Table No. 3: Visual acuity. (n=280)


No of patients on presentation	Visual acuity	No of patients on final presentation	Visual acuity
90 (32.14%)	6/6	200 (71.43%)	6/6
130 (46.43%)	6/9	46 (16.43%)	6/9
12 (4.29%)	6/18	04 (1.43%)	6/18
18 (6.43%)	6/36	07 (2.5%)	6/36
23 (8.21%)	6/24	17 (6.07%)	6/24
05(1.79%)	<6/60	04 (1.43%)	<6/60
02 (0.71%)	PI+ve	02 (0.71%)	PI+ve
Total: 280 (100%)	-----	280 (100%)	-----

Table No.4: Distribution manner of injury. (n=280)

Manner of injury	No	%age
Accidental	212	75.72
Homicidal	68	24.28
Total	280	100%



Graph No. 1 Distribution of ocular injuries (n=27542)



Graph No. 2: Distribution of gender (n=280)

DISCUSSION

Injuries to the eye are more common & deleterious in their effects specially in developing countries than developed. In this study among 280 patients 250 (89.29%) were male and female 30 (10.71%) with M:F ratio of 8.34:1 while in other studies M:F ratios 2.56:1⁵, 2.85:1⁶, 3:1⁷, 4.78:1⁸,5:1⁹,5.49:1¹⁰.High

incidence of ocular injury was seen in males because of occupational exposure, taking part in aggressive activities. Our study shows the age distribution for the occurrence of ocular trauma is maximum 106 (37.86%) in 21-30 years age group and the second peak 82 (29.28%) in the 11-20 years. The propensity towards young and school going children in our study is respectively parallel to the trends reported by Akram et al¹¹, Gyasi et al¹², Maurya et al¹³, Sukati & hansraj^{14,15}. Higher preponderance in children & young individuals indicates that children have decreased ability to detect and avoid potential hazards because of non-supervision at play & domestic activities while young ones are more aggressive, occupational exposure, participation in dangerous sports & hobbies, alcohol use & risk taking behavior. Our study represents 71.43% injuries caused by sharp objects and studies conducted by L.O One we 13.4%¹⁶ & Guly et al 3.3%¹⁷ are in contrast. Injuries by blunt weapon in our study shows 17.86% that is in consistent to study conducted by Sintuwong & Winitchai 19.2%⁸ and in contrast in studies by Voon et al 12.6%¹⁸, Nqo & Leo 4%¹⁹, Gyasi et al 41.3%¹², Zelalem Addisu 40%²⁰ & Zaqeiboumet al 60%²¹. Foreign bodies lodged in eyes in our study is 10.71% while in studies by Babar et al 11.8%²², Cilino et al 16.8%¹⁰, Aghadost & others 6.1%⁹ are almost in same incidence. Higher incidence by sharp objects is due to urban population, industrial area where there is easy accessibility to such objects. This study shows a good initial visual acuity 78.57% study by Cilino et al¹⁰ because of minor injuries in most of the cases involving the eye- lid, conjunctiva, cornea and anterior chamber of an eye having association with a good visual prognosis than posterior chamber trauma and the modern surgical & medical management as well as the better awareness of urban population than rural population to reach the Eye emergency room earlier. Our study shows in 212 (75.72%) cases accidental injuries which is in parallel to study conducted by Vats & other authors 87.1%²³ and assault cases are 68 (24.28%) in our study that is nearly in consistent with study of Babar et al 37.37%²⁴ and in contrast to Shashikal & others 3.3%²⁵. The high rate of accidental injuries is due to ill-fitting or non-availability of protective measures at work place, poor vision due to fogging from sweat, specially in children & young people because of decreased ability to detect & avoid potential hazards, unsupervision at play and domestic activities as well as taking part in dangerous sports.

CONCLUSION

The mostly cases were of male gender in the age of children & adults. The injuries caused mainly by sharp objects resulted in visual impairment initially but with good final vision because of advanced technology. The accident was the main manner of injury.

Recommendation: Adequate supervision of children Educational measures to reduce accidents by arranging seminars, information through media etc.

Maintain Eye Trauma Registry in the department of ophthalmology nationwide in order to provide proper ocular trauma statistics.

Acknowledgement: The authors are thankful to Mr. Nadeem Ali Khokhar, Computer Operator for typing the manuscript.

REFERENCES

1. Durani J. Accidental Eye Injuries. Pak J Ophthalmol 1997;13(4):96-7.
2. Caroline. Ocular Injuries. J R Coll Surg Edinb 1999;44(10): 317-23.
3. Ahmed I, Seema N, Humayoon M, Raja A. Weapon of offence used in bodily Medico-legal Injuries in a Rural Area. Med Forum 2011;22(5):51-53.
4. Kirshan V, editor. Injuries medicolegal considerations and types. Text book of Forensic Medicine and Toxicology. 3rd ed. New Delhi: Elsevier Publisher; 2005.p. 300.
5. Beby F, Kodjikian L, Roche O, Donate D, Kouassi N, Burillion C, Denis P. Perforating ocular injuries in children: a retrospective study of 57 cases. *Francais D'ophtalmologie* 2006; 29(1): 20-23.
6. Pandita A, Merriman M. Ocular trauma epidemiology: 10-year retrospective study. NZMA 2012; 125(1348):1/7-7/7.
7. Mc Call BP, Herwitz LB, Taylor DA. Occupational eye injury and risk reduction; Kentucky worker's Compensation analysis 1994-2003. Inj Prev 2009; 15: 176-182.
8. Sintuwong S, Winitchai R. Predictive factors of visual outcome in open globe injuries in Thailand: a prospective study. Asian Biomedicine 2011;5 (2): 289-94.
9. Aghadoost D, Fazel MR, Aghadoost HR. Pattern of Paediatric ocular trauma in Kashan. Archives of Trauma Res 2012;1(1): 135-7.
10. Cilino S, Casuccio A, Pace FD, Pillitteri F, Cilino G. A five-year retrospective study of the epidemiological characteristics and visual outcomes of patients hospitalized for ocular trauma in a Mediterranean area. BMC Ophthalmology 2008; 8 (6): 121-9.
11. Shahwani MA, Hameed K, Jamali B. Ocular Injuries: Its Etiology and Consequences in Balochistan. Pak J Ophthalmol 2006; 22 (2): 82-84.

12. ME Gyasi, WMK Amontku, MA Aduik. Epidemiology of hospitalized ocular injuries in the upper East region of Ghana. Ghana Med 2007; 41(4): 171-75.
13. Maurya RP, Sinhak, Den PR, Singh VP, Singh MK, Bhushan P. A clinico-Epidemiological study of ocular trauma in Indian University students. Pak J Ophthalmol 2013; 29(2): 80-88.
14. Sukati VN, Hansraj R. Retrospective analysis of eye injuries in rural Kwazulu-Natal, South Africa. S Afr Optom 2012; 71(4): 159-65.
15. Sukati VN, Hans raj Characteristics of injuries in urban Kwazulu-Natal Province, South Africa. S Afr Optom 2013; 72(3):119-26.
16. Onyekwe LO. Spectrum of eye injuries in children in Guinness Eye Hospital, Onitsha. The Nigerian J Surg Res 2001; 3(3): 126-32.
17. Guly CM, Guly HR, Lecky FE. Ocular injuries in patients with major trauma. Emerg Med J 2008; 23(2): 915-17.
18. Voon LW, See J, Wong TY. Epidemiology of ocular Trauma in Singapore, perspective from the emergency service of large Tertiary Hospital. Eye 2001; 15(1): 75-81.
19. Nqo CS, Leo SW. Industrial accident-related ocular emergencies in a tertiary Hospital Singapore. Singapore Med J 2008; 49(4):280-285.
20. Addisu Z. Pattern of ocular trauma in rural Ethiopia, in Garbat Eye Hospital. Central Ethiopia. Ethiop J Health Dev 2011; 25(2): 150-55.
21. Zaqeiboum BM, Tostanoski JR, Kerner DJ, Hersh PS. Urban eye trauma, a one year prospective study. Ophthalmol 1993; 100(6):851-60.
22. Babar TF, Jan S, Gul L, Tariq M, Khan MU, Zaman M, Khan MD. Pattern of pediatric ocular trauma in Hayatabad Medical Complex Peshawar. Pak J Med 2006; 45(1): 145-9.
23. Vats S, Murthy GVS, Gogoi M. Ind J Ophthalmol 2008; 58(4): 313-18.
24. Babar TE, Khan MT, Moawal MZ, Shah SA, Murad Y, Khan MD. Patterns of ocular trauma. J Coll Physicians Surg Pak 2007; 17(3): 148-53.
25. Shashikala P, Sadiqulla M, Shivakumar D, Prakash KH. Profile of ocular trauma in industries-related Hospital. Ind J occup. Environ Med 2013;17(2): 66-70.

Address for Corresponding Author:

Dr. Muhammad Afzal Pechuho,

Assoc. Prof. & HOD Ophthalmology,

Ghulam Mohammad Mahar Medical College,

Sakur

Cell No.: 0300-3423761

Electronic Copy