

Hysterosalpingography in Uterine and Tubal Factor Subfertility - An Outdated Procedure or a Reemergent Study?

Hysterosalpingography
in Uterine and Tubal
Factor Subfertility

Fauzia Afridi¹, Arzoo Gul Bangash², Aysha Afridi³ and Zainab Afridi⁴

ABSTRACT

Objective: to determine frequency of various abnormalities detected on hysterosalpingography in patients presenting with subfertility.

Study Design: Descriptive / cross-sectional study

Place and Duration of Study: This study was conducted at the Department of Obstet. and Gynae, Noshera Medical College, Noshera. from October 2013 to March 2014.

Materials and Methods: This study involved 318 patients presenting to outpatient Gynae clinic of Northwest General Hospital and Research Hospital, Peshawar with primary or secondary infertility of at least one year. After initial history, examination and consent, HSG was performed using set criteria and the films reported by consultant radiologist.

Results: Mean age of the patients was 29.6 years with 25-36 years being the commonest age group. 58.5% patients had primary infertility whereas 41.5% patients had secondary infertility. Majority of the patients (84%) had normal HSG whereas only 16% showed some abnormality. The abnormalities detected on HSG included Tubal block (13.8%...unilateral:9.4%; bilateral:4.4%), Hydrosalpinx (2.5%...unilateral:2.5%; bilateral:0.6%), Peritubal adhesions(1.6%) and uterine abnormalities (2.8%). The most frequent uterine abnormality was bicornuate uterus (1.6%) which was more common in primary infertility patients, followed by submucous fibroid/ polyp (0.6%). Proximal tubal block was more common than distal tubal block.

Conclusion: The majority of patients in this study had normal HSG and most of these can be spared laparoscopy, given the high specificity of HSG for tubal patency. Therefore, HSG is still the 1st line investigation in the workup of tubal factor for infertility.

Key Words: Infertility, Hysterosalpingography, tubal block.

Citation of article: Afridi F, Bangash AG, Afridi A, Afridi Z. Hysterosalpingography in Uterine and Tubal Factor Subfertility - An Outdated Procedure or a Reemergent Study? Med Forum 2017;28(6):91-95.

INTRODUCTION

The desire to procreate is intrinsic to every woman. Inability to do so leads to unimaginable physical, psychological, economic and social distress.^{1,2,3} Infertility is defined as the inability of a couple to conceive following 12 months or more of regular unprotected intercourse.^{4,5,6} According to WHO, 1 in every 4 couples in developing countries are affected by infertility.⁷ However, its incidence ranges from 6.6-32.6%⁸, depending on the duration of infertility used in definition and the population studied, e.g., community or hospital clinics.⁹

The causes of infertility include anovulation (20%), tubal factor (11-30%), male factor (20-30%).¹⁰ Initial workup of infertile couple should include Semen analysis (analysis of male factor), transvaginal ultrasound scan (pelvic anatomy), follicular phase gonadotrophins and mid-luteal progesterone (ovarian reserve and ovulation).^{11,12}

Routine tubal patency testing, being invasive, is debatable in the infertility workup.¹² It should be offered after taking into account the overall treatment needs of the couple. The most commonly used tests of tubal patency include Hysterosalpingography (HSG), Laparoscopy and Dye test and Hysterocontrast sono-salpingography.^{13,14} Other less commonly used tests include Selective Salpingography and tubal catheterization, transvaginal hydrolaparoscopy, fertiloscopy, salpingoscopy, fallopscopy and chlamydia antibody testing.^{15,16,18,19}

Laparoscopy is widely considered to be the gold-standard for assessing tubal patency, as it enables direct visualization of pelvis and allows treatment of mild endometriosis and periadnexal adhesions.²⁰ However, it is more invasive, requires general anesthesia and carries risk of injury to abdominal organs.²¹ In contrast, HSG being cheap, widely available, less invasive and an outpatient procedure, can be regarded as a better 1st line investigation for assessing tubal patency. It has a high specificity of 83%, making it a better test for

¹. Department of Obstet & Gynae, Nowshera Medical College, Nowshera.

². Department of Obstet & Gynae, Mardan Medical Complex, Mardan.

³. Department of Radiology, Maqsood Medical Complex, Peshawar.

⁴. Department of Radiology, Institute of Kidney Diseases, Peshawar.

Correspondence: Dr. Fauzia Afridi, Assistant Professor, Nowshera Medical College, Nowshera.

Contact No: 0333-9154231

Email: Fauzia_afridi@yahoo.com

identification of tubal patency. Its limitations include failed catheterization or instrumentation., and false positives (sensitivity 65%) due to tubal spasm or debris.^{22,23,24}

HSG is a several decades old investigation now. The improvements in radiology/ fluoroscopic services, in addition to the advent of several new techniques for assessing tubal pathology has necessitated the need to re-appraise its role in the current era of ART. This study was conducted with the aim to evaluate its role in this context.

MATERIALS AND METHODS

This descriptive study was carried out at Northwest General Hospital and Research Center, Peshawar from October 2013 to March 2014. The subjects were 318 patients, selected by non-purposive convenience sampling, presenting to outpatient Gynae clinic, with primary or secondary subfertility of at least 1 year. Those patients whose husbands were abroad for the last one year or had abnormal semen parameters were excluded from the study. The procedure and purpose of the study was explained to the subjects and informed

consent was taken. Subjects were given an intramuscular injection of Diclofenac Sodium 15 minutes before the procedure. With patient in dorsal position, uterine size and position were assessed first digitally and then with uterine sound. After introducing Cuscos speculum and holding anterior cervical lip with tenaculum, size 8 F catheter was introduced through cervix and retained. In Radiology department, 10-20 ml of radio-opaque dye (Urograffin) was introduced through the catheter under fluoroscopic control and X-ray films taken. Antibiotic prophylaxis was given to all patients after the procedure.the findings on HSG films were reported by consultant radiologist.

All the relevant data was entered on to a proforma and analysed using SPSS 20. Results were presented as graphs and tables.

RESULTS

This study involves 318 patients presenting with infertility. The ages of the patients ranged from 17-45 years. Mean age was 29.64+/_5.32 SD. The results are tabulated as below:

Table No.1: Age Groups

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	< 20 yrs	9	2.8	2.8	2.8
	20-25	67	21.1	21.1	23.9
	26-30	113	35.5	35.5	59.4
	31-35	77	24.2	24.2	83.6
	36-40	46	14.5	14.5	98.1
	> 40	6	1.9	1.9	100.0
	Total	318	100.0	100.0	

Table No.2: Type of Infertility

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	primary	186	58.5	58.5	58.5
	secondary	132	41.5	41.5	100.0
	Total	318	100.0	100.0	

Table No.3: HSG findings

		Frequency	Percent	Valid Percent	Cumulative Percent
Normal		267	84.0	84.0	84.0
Abnormal		51	16.0	16.0	100.0
Total		318	100.0	100.0	

Table No.4: Tubal Block Laterality

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	0	274	86.2	86.2	86.2
	unilateral	30	9.4	9.4	95.6
	bilateral	14	4.4	4.4	100.0
	Total	318	100.0	100.0	

Table No.5 :Hydrosalpinx

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	0	310	97.5	97.5	97.5
	unilateral	6	1.9	1.9	99.4
	bilateral	2	.6	.6	100.0
	Total	318	100.0	100.0	

Table No.6: Peritubal Adhesion

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	0	313	98.4	98.4	98.4
	1	5	1.6	1.6	100.0
	Total	318	100.0	100.0	

Table No.7: Uterine Abnormalities

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	0	309	97.2	97.2	97.2
	uterus didelphys	1	.3	.3	97.5
	bicornuate uterus	5	1.6	1.6	99.1
	infantile uterus	1	.3	.3	99.4
	submucous fibroid/polyp	2	.6	.6	100.0
	Total	318	100.0	100.0	

Table No.8: Tubal Block Site

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	0	274	86.2	86.2	86.2
	proximal	26	8.2	8.2	94.3
	distal	18	5.7	5.7	100.0
	Total	318	100.0	100.0	

DISCUSSION

This study involves 318 women presenting with subfertility. Most of the patients (35.5%) were in the age group / of 26-30 years. This is in accordance with studies conducted in other developing and under-developed countries,²⁵ where early marriage and child bearing is a norm, but in contrast to studies in the developed world²⁶ where there is trend to delaying marriage and child birth, hence infertile couple present later in age.

In this study, Primary infertility was the major type of infertility, comprising of 58.5% of patients whereas 41.5% of patients presented with secondary infertility. Studies on infertility have shown a similar trend in that secondary infertility ,although less frequent than primary infertility , is on the rise.^{7, 27} This is probably due to increasing rates of caesarian sections and Pelvic inflammatory disease, resulting in tubal block and also due to increasing age of patients and prolonged birth-spacing resulting in diminishing ovarian reserve with time.

Regarding HSG findings, 84% of subjects had normal HSG findings whereas 16% had one or more abnormal findings on HSG. Studies on HSG conducted so far report wide variations in the proportions of abnormal HSG.^{25,26,27} This depends on whether patients have been referred after initial workup elsewhere and also on the prevalence of PID in that set-up; both situations leading to increased rates of abnormal HSG as much as 50%. In our study, the yield of HSG was low (only 16% abnormal HSG). This could be either because the indication was too broad or there was a selection bias in that those who were considered to have pelvic

pathology like PID, endometriosis, adhesions were directly referred for laparoscopy.²⁶

The commonest abnormality detected on HSG was Tubal block (13.8%.....unilateral:9.4%; bilateral: 4.4%). Similar results were obtained in a study conducted in Oman by Subhi et al.²⁸ Hydrosalpinx accounted for 2.5% cases, out of which unilateral were 1.9% and bilateral were 0.6%. The diagnosis of hydrosalpinx is significant in that preventive salpingectomy is recommended in such patients prior to IVF to improve implantation rates.²⁹

Peritubal adhesions, detected by convoluted fallopian tubes and/or loculation of spillage of contrast medium in the peritoneal cavity,³⁰ were seen in 1.6% patients. Although HSG can detect peritubal adhesions in upto 75% cases,³⁰ laparoscopy is the gold-standard for their diagnosis and HsG should not be used for this purpose²².

Uterine abnormalities were detected in 2.8% patients. The most frequent uterine abnormality was bicornuate uterus(1.6%) which was more common in primary infertility patients, followed by submucous fibroid/polyp(0.6%) which was more common in secondary infertility patients. Abnormalities in the uterine cavity detected on HSG should be further evaluated by hysteroscopy.

Determination of tubal block site is important as it can help us in choosing the type and determining prognosis of tubal reconstructive surgery.³¹ In this study, 8.2% of patients had proximal tubal block as evident by the lack of tubal filling on HSG whereas 5.7% of patients had distal tubal block in which the tube was outlined however there was no spill of the dye. A similar result was obtained by a study conducted in Nepal by Poonam.²⁷

In this study, majority of patients (84%) had normal HSG. Studies have revealed that , in infertile couples having normal HSG, there is a four-fold increase in pregnancy rate during first three months after HSG and, although not by the same proportion, but the trend remains so up to one year after HSG.it is in this subset of patients who can be spared initially from invasive investigations like laparoscopy and hysteroscopy. Only those patients can be referred for laparoscopy who fail to conceive within 3-12 months of HSG. After 12 months, there is no difference in the rate of pregnancy among those who had and and those who didn't have HSG. By following such criteria for laparoscopy, the chance of finding a pelvic pathology, and/or the need for change in treatment plan after laparoscopy is increased, hence making it more cost effective. Thus, HSG can not only short-list patients for laparoscopy,³²it may have a therapeutic effect on the remaining patients.³³.

CONCLUSION

HSG is a valuable initial investigation in the work up of infertility, even in the modern era of ART. It is fairly reliable in establishing patency of tubes. However, the small minority of patients with blocked tubes or intra-cavitary pathology need to be evaluated further by hysterosalpingography and hysteroscopy respectively..

Conflict of Interest: The study has no conflict of interest to declare by any author.

REFERENCES

- Ezzel W. The impact of Infertility on Women's Mental Health. *N C Med J* 2016;77(6):427-428.
- Abbey A. adjusting to infertility. In:Harvey JD, Miller ED, editors. *Loss and Trauma: General and Close Relationship Perspectives*. Ann Arbor MI: Edward Brothers; 2000.
- Bhatti LI, Fikree FF, Khan A. The quest of infertile women in squatter settlements of Karachi, Pakistan: a qualitative study. *Social Sci Med* 1999;49(5):637-49.
- Zegers-Hochschild F, Adamson GD, de Mouzen J, Ishihara O, Mansour R, Nygren K et al. The International Committee for Monitoring Assisted Reproductive Technology (ICMART) and the World Health Organization (WHO) Revised Glossary on ART Terminology, 2009. *Hum Repr* 2009;24(11):2683-2687.
- Practice Committee of the American Society for Reproductive Medicine; Definitions of infertility and recurrent pregnancy loss. *Fertil Ster* 2008;90(5).
- National Institute for Health and Care Excellence. *Fertility problems: assessment and treatment*. London: NICE; 2013.
- Mascarenhas MN, Flaxman SR,Boerma T, Vanderpoel S, Stevens GA. National, Regional and Global Trends in Infertility prevalence since 1990: A systematic analysis of 277 Health Surveys. *PLoS Med* 2012;9(12):e1001356.
- Mark Hamilton. *Infertility*. In:Keith Edmonds D ed. *Dewhurst's Textbook of Obstetrics and Gynaecology*, 8th edition.UK. Blackwell Publishing Ltd; 2012.
- Gurunath S, Pandian Z, Anderson RA, Bhattacharya S. Defining infertility_ a systematic review of prevalence studies. *Hum Repr Update* (2011)17(5):575-588
- Hull MGR, Glazener CMA, Kelly NJ, et al. Population study of causes, treatment and outcome of infertility. *BMJ* 1985;291:1693-1697.
- World Health Organization. *WHO Manual for the standardized investigation and diagnosis of the infertile couple*. Cambridge, UK: Cambridge University Press; 1993.
- Forti G,Krausz C. Evaluation and Treatment of the Infertile Couple. *J Clin Endocrinol Metab* 1998; 83(12):4177-4188.
- Dun EC, Nezhat CH. Tubal factor infertility: diagnosis and management in the era of assisted reproductive technology.*Obstet Gynecol Clin North Am* 2012;39(4):551-66.
- Phillips CH, Benson CB, Ginsburg ES,Frates MC.Comparison of uterine and tubal pathology identified by transvaginal sonography, hysterosal pingography, and hysteroscopy in female patients with infertility. *Fertil Res Pract* 2015;1:20.
- Papaioannou S,Afnan M,Girling AJ,Coomarasamy A,McHugo JM,Sharif K.The potential value of tubal perfusion pressures measured during selective salpingography in predicting fertility. *Hum Reprod* 2003;18(2):358-363.
- Thomas K, Coughlin L, Mannion PT, Haddad NG. The value of Chlamydia trachomatis antibody testing as part of routine infertility investigations. *Hum Reprod* 2000;15(5):1079-1082.
- Papaioannou S, Bourdrez P, Varma R,Afnan M,Mol BWJ, Coomarasamy A.Tubal evaluation in the investigation of subfertility:A structured comparison of tests. *Br J Obstet Gynaecol* 2004;111(12):1313-1321.
- WatreLOT A,Dreyfus JM,Andine JP.Evaluation of the performance of fertiloscopy in 160 consecutive infertile patients with no obvious pathology. *Hum Reprod* 1999;14(3):707-711.
- Maarana R, et al. *Curr Opin Obstet Gynecol* 1996;8(4):257-260.
- Rowe PJ, Comhaire FH,Hargreave TB,Mahmoud AMA. *WHO manual for the standardized*

- investigation of the infertile couple. Cambridge UK: Cambridge University Press; 1993.
21. Tanahatoe SJ, Hompes PGA, Lambalk CB. Investigation of the infertile couple: should diagnostic laparoscopy be performed in the infertility workup programme in patients undergoing intrauterine insemination? *Hum Reprod* 2003;18(1):8-11.
 22. Swart P, Mol BWJ, van der Veen F, van Beuden M, Redekop WK, Bossuyt PMM. The accuracy of hysterosalpingography in the diagnosis of tubal pathology: a meta-analysis. *Fertil Steril* 1995; 64(3):486-491.
 23. The ESHRE Capri Workshop Group. Optimal use of infertility diagnostic tests and treatments. *Hum Reprod* 2000;15(3):723-732.
 24. Lim CP, Hasafa Z, Bhattacharya S, Maheshwari A. Should a hysterosalpingogram be a first-line investigation to diagnose female tubal subfertility in the modern subfertility workup? *Hum Reprod* 2011;26(5):967-971.
 25. Kiguli-Malwadde E, Byanyima RK. Structural findings at hysterosalpingography in patients with infertility at two private clinics in Kampala. Uganda. *Afr Health Sci* 2004;4(3):178-181.
 26. Schankath AC, Fasching N, Urech-Ruh C, Hohl MR, Kubik-Huch RA. Hysterosalpingography in the workup of female infertility: indications, technique and diagnostic findings. *Insights Imaging* 2012;3(5):475-483.
 27. Poonam. The role of hysterosalpingography in cases of subfertility. *Kathmandu Univ Med J* 2007;5(4):456-460.
 28. Subhi TA, Jashnmi RNA, Khaduri MA, Gowri V. Prevalence of Tubal Obstruction in the Hysterosalpingogram of women with primary and secondary infertility. *J Reprod Infertil*. 2013; 14(4):214-216.
 29. Strandell A, Lindhard A, Waldenström U, Thorburn J, Janson PO, Hamberger L. Hydrosalpinx and IVF outcome: a prospective, randomized multicenter trial in Scandinavia on salpingectomy prior to IVF. *Hum Reprod* 1999;14(11):2762-2769.
 30. Karasick S, Goldfarb AF. Peritubal adhesions in infertile women: Diagnosis with Hysterosalpingography. *AJR* 1989;152:777-779.
 31. Suresh YN, Narvekar NN. The role of tubal patency tests and tubal surgery in the era of assisted reproductive techniques. *Obstet Gynaecol* 2014;16(1):37-45.
 32. Berker B, Sukur YE, Aytac R, Atabekoglu CM, Sonmezer M, Ozmen B. Infertility work-up: To what degree does laparoscopy change the management strategy based on hysterosalpingography findings? *J Obstet Gynecol Res* 2015;41(11):1785-1790.
 33. Cundiff G, Carr BR, Marshburn PB. Infertile couples with a normal hysterosalpingogram: Reproductive outcome and its relationship to clinical and laparoscopic findings. *The J Reprod Med* 1995;40(1):19-24.