

Role of Adrenaline Sclerotherapy for Controlling Upper GI Bleeding

Muhammad Waqas¹, Imdad Ali Ansari³, Mujtaba Jaffary⁴ and Ghias-un-Nabi Tayyab²

ABSTRACT

Objectives: To determine the efficacy of adrenaline sclerotherapy to control upper GI bleeding

Study Design: Observational / descriptive study.

Place and Duration of Study: The study was conducted at Gastroenterology/Medical Ward, Lahore General Hospital, Lahore in collaboration with Chandka Medical College Hospital, Larkana from 1st January 2016 to 30th June 2016.

Materials and Methods: In this study a total of 100 cases of either gender between 20-60 years of age with upper GI bleed within last 24 hours were enrolled. The basic demographic data including age, gender, clinical presentation were recorded. Clinical efficacy of adrenaline was considered if there was no further upper GI bleeding upto 72 hour after endoscopy.

Results: The mean age was recorded as 38.41±11.83 years, 57% (n=57) were males and 43% (n=43) were females, efficacy of adrenaline sclerotherapy in controlling upper GI bleeding in was recorded in 66% (n=66) of the cases.

Conclusions: The efficacy of adrenaline sclerotherapy is significantly higher and encouraging for controlling upper GI bleeding and this modality may be used for effective management of upper GI bleeding.

Keywords: Upper GI bleeding, Cancer population, Adrenaline sclerotherapy, Efficacy

Citation of article: Waqas M, Ansari IA, Jaffary M, Tayyab GN. Role of Adrenaline Sclerotherapy for Controlling Upper GI Bleeding. Med Forum 2017;28(5):163-165.

INTRODUCTION

Upper gastrointestinal (GI) bleeding is known as bleeding source proximal to the ligament of Treitz. Various upper GI bleeding cases may cause iron-deficiency anemia but they may not generally present in emergency case. Upper GI bleeding cases are characterized by melena, hematemesis, hematochezia and evidence of compromise for hemodynamic e.g. shock, dizziness and syncope episodes. Usually they are caused by major haemorrhage from ulcers, Dieulafoy lesions, varices, Mallory-Weiss tears; and neoplasms. These cases are required to be admitted to Intensive Care Unit and urgently a consultant gastroenterologist may be approached. The cases presenting with massive bleeding should be observed for surgery.¹⁻⁵ Upper endoscopy is a choice as diagnostic modality while it is also considered as a choice of treatment for acute upper GI bleeding.^{6,7}

Being a potentially life-threatening condition, it needs active and appropriate management. Despite of advancement in therapeutic management, the rate of mortality has not changed i.e. 10-14% and relates to the longer life expectancy along with increased number of co-morbidities.⁸

Epinephrine is known as the popular injectate being used for injection therapy and promotes hemostasis by using local tamponade.⁹ It arrests around 80% of the bleeding.¹⁰ Initially, it reduces markedly or stops the bleeding but the bleeding may recur more than or equal to 20 minutes after therapy if the residual clot is not already formed as the epinephrine injection is absorbed and flushed into the circulation and however, its local effect, wears off.¹¹ It is the perfect initial agent as it is an easy to administer and generally reduces the rate of bleeding thereby. It is a common method of achieving hemostasis owing to its low risk, low cost and easy availability but local data is very limited assessing the efficacy of the endoscopic adrenaline use in patients presenting with upper GI bleed. However, the current trial was designed to assess the effectiveness of adrenaline for Upper GI bleeding.

MATERIALS AND METHODS

In this study a total of 100 cases of either gender between 20-60 years of age with upper GI bleed within last 24 hours were enrolled. The study was conducted at Gastroenterology/Medical Ward, Lahore General Hospital, Lahore in collaboration with Chandka Medical College Hospital, Larkana from 1st January 2016 to 30th June 2016. Informed consent of the

¹. Department of Medicine / Gastroenterology², Lahore General Hospital/Postgraduate Medical Institute, Lahore.

². Department of Medicine, CMC, SMBBU, Larkana

¹. Department of Medicine, Continental Medical College Lahore

Correspondence: Dr. Muhammad Waqas, ¹Senior Registrar, Department of Medicine Unit-1, Lahore General Hospital/Postgraduate Medical Institute Lahore.

Contact No: 0321-6660213

Email: drwaqas@live.com

Received: March 11, 2017;

Accepted: April 20, 2017

patients was obtained to include their data for publication. The basic demographic data including age, gender, clinical presentation were recorded. After a detailed history and physical examination, 5 to 10ml of diluted (1:10,000) adrenaline was injected endoscopically around bleeding point by dedicated gastroenterologist. Clinical efficacy of adrenaline was considered if there was no further UGIB upto 72 hour after endoscopy. The data was analyzed for clinical efficacy by using simple frequency and percentage.

RESULTS

Age distribution of the patients was done showing that 55% (n=55) were between 12-40 years of age while 45% (n=55) were between 41-70 years of age, mean±SD was calculated as 38.74±11.52 years (Table 1). Patients were distributed according to gender, it shows 57% (n=57) were male and 43% (n=43) were females (Table 2). The efficacy of adrenaline sclerotherapy in controlling upper GI bleeding was recorded in 65% (n=65) of the cases (Table 3)

Table No.1: Age distribution (n=100)

| Age (years) | No. | % |
|-------------|-----|------|
| 12 – 40 | 55 | 55.0 |
| 41 - 70 | 45 | 45.0 |

Table No.2: Gender Distribution (n=100)

| Gender | No. | % |
|--------|-----|------|
| Male | 57 | 57.0 |
| Female | 43 | 43.0 |

Table No.3: Efficacy of adrenaline sclerotherapy in controlling upper GI bleeding (n=100)

| Efficacy | No. | % |
|----------|-----|------|
| Yes | 65 | 65.0 |
| No | 35 | 35.0 |

DISCUSSION

Our data regarding efficacy of adrenaline sclerotherapy for controlling upper GI bleed is in agreement with previous data showing that Argon plasma coagulation (APC), injection sclerotherapy using 1:10,000 adrenaline, endo-clips and heat probe effectively controlled the bleeding in 75.9,¹¹ 75,¹² 87¹³ and 97¹⁴ percent of patients respectively, the findings of these studies support our results. However, some of the above studies showing higher efficacy than ours.

Another study by Chung et al¹⁵ revealed that injection with 1:10000 adrenaline alone reduces active bleeding in majority of the cases, however, rebleed rates was not recorded in his study. Wardehoff and Gros¹⁶ revealed that injection of the sclerosant polidocanol decreases the risk of rebleed in cases presenting with major stigmata of recent haemorrhage, while the efficacy of

sclerosants alone as a management for active bleeding is not well cleared.

A prospective trial including 78 cases with active bleeding or non-bleed visible vessel from a peptic ulcer (PU) who administered with a mean of 16.5 ml of epinephrine, 12 cases (15.4%) had recur bleed.¹⁷

In absence of no strict criteria regarding the quantity, a total of higher volume¹⁸, about 30 mL¹⁹ may be optimized as hemostasis and associated with reduced rate of rebleeding.^{19,20}

However, a second injection is not generally applied considering the fact that it does not confer more advantage over epinephrine injection alone, but it may can potentiate epinephrine-induced mucosal injury.^{15,21}

Though we found a higher rate of efficacy of adrenaline sclerotherapy to control upper GI bleed but these findings are primary in our setup and needs some-other trials to validate this data.

CONCLUSION

The efficacy of adrenaline sclerotherapy is significantly higher and encouraging for controlling upper GI bleeding and this modality may be used for effective management of upper GI bleeding.

Conflict of Interest: The study has no conflict of interest to declare by any author.

REFERENCES

- Laine L, Yang H, Chang SC, Datto C. Trends for incidence of hospitalization and death due to GI complications in the United States from 2001 to 2009. *Am J Gastroenterol* 2012;107:1190–5.
- Bhutta S, Jamil M, Aziz K, Wasimuddin. An etiological study of uppergastrointestinal bleeding; *J JRM* 2012;16(1):31-3.
- Shebani S, Kim JJ, Chen B, Park S, Saberio S, Keyashian K, et al. Natural history of acute upper GI bleeding due to tumours:short-term success and long-term recurrence with or without endoscopic therapy. *Aliment Pharmacol Ther* 2013;38:144–50.
- Yarris JP, Warden CR. Gastrointestinal bleeding in cancer patient. *Emerg Med Clin North Am* 2009; 27(3):363-79.
- Maluf-Filho F, daCosta-Martins B, deLima MS, Leonardo VD, Retes FA, Kawaguti FS, et al. Etiology, endoscopic management and mortality of upper gastrointestinal bleeding in patients with cancer. *United Eur Gastroenterol J* 2013;1:60.
- Charatcharoenwitthaya P, Pausawasdi N, Laosanguaneak N, Bubthamala J, Tanwadee T, Leelakusolvong S. Characteristics and outcomes of acute upper gastrointestinal bleeding after therapeutic endoscopy in the elderly. *World J Gastroenterol* 2011; 17(32):3724-32.

7. Adang RP, Vismans JF, Talmon JL, Hasman A, Ambergen AW, Stockbrügger RW. Appropriateness of indications for diagnostic upper gastrointestinal endoscopy: association with relevant endoscopic disease. *Gastrointest Endosc* 1995;42(5):390.
8. Savides TJ, Jensen DM, Cohen J. Severe upper gastrointestinal tumor bleeding endoscopic findings, treatment, and outcomes. *Endoscop* 1996;28:244-8.
9. Simons FE, World Allergy Organization. Epinephrine auto-injectors: first-aid treatment still out of reach for many at risk of anaphylaxis in the community. *Ann Allergy Asthma Immunol* 2009;102:403.
10. Calvet X, Vergara M, Brullet E, Gisbert J P& Campo R. Addition of a second endoscopic treatment following epinephrine injection improves outcome in high-risk bleeding ulcers. *Gastroenterol* 2004;126:441-50.
11. Cappell MS, Friedel D. Acute nonvariceal upper gastrointestinal bleeding: endoscopic diagnosis and therapy. *Med Clin North Am* 2008;92:511-50.
12. Kanai M, Hamada A, Endo Y, Takeda Y, Yamakawa M, Nishikawa H. Efficacy of argon plasma coagulation in nonvariceal upper gastrointestinal bleeding. *Endoscop* 2004;36(12(e.g. 2)):1085-8.
13. Sung JJ, Tsoi KK, Lai LH, Wu JC, Lau JY. Endoscopic clipping versus injection and thermocoagulation in the treatment of non-variceal upper gastrointestinal bleeding: A meta-analysis. *Gut* 2007;56:1364-73.
14. Church NI, Dallal HJ, Masson J. A randomised trial comparing heater probe plus thrombin with heater probe plus placebo for bleeding peptic ulcer. *Gastroenterol* 2003;125:396-403.
15. Chung SCS, Leung JWC, Steele RJC, Crofts TJ, Li AKC. Endoscopic injection of adrenaline for actively bleeding ulcers: a randomised trial. *BMJ* 1988;2:1631-3.
16. Wordehoff D, Gros H. Endoscopic haemostasis by injection therapy in high-risk patients. *Endoscop* 1982;14:196-9.
17. Lin HJ. A prospective, randomized trial of large-versus small-volume endoscopic injection of epinephrine for peptic ulcer bleeding. *Gastrointest Endosc* 2002;55:615-9.
18. Park CH, Lee SJ, Park JH, Lee WS, Joo YE, Kim HS, et al. Optimal injection volume of epinephrine for endoscopic prevention of recurrent peptic ulcer bleeding. *Gastrointest Endosc* 2004; 60(6): 875-80
19. Jensen DM, Machado GA. Endoscopic hemostasis of ulcer hemorrhage with injection, thermal, and combination methods. *Tech Gastrointest Endosc* 2005;7:124-31
20. Laine L, Long GL, Bakos GJ, Vakharia OJ, Cunningham C. Optimizing bipolar electrocoagulation for endoscopic hemostasis: assessment of factors influencing energy delivery and coagulation. *Gastrointest Endosc* 2008; 67(3): 522-8
21. Kovacs TO. Management of upper gastrointestinal bleeding. *Curr Gastroenterol Rep* 2008;10:535-42.

Electronic Copy